

## National checklist for aquatic alien species in Germany

Stephan Gollasch<sup>1</sup> and Stefan Nehring<sup>2</sup>

<sup>1</sup>*GoConsult, Bahrenfelder Str. 73a, 22765 Hamburg, Germany*

*E-mail: [sgollasch@aol.com](mailto:sgollasch@aol.com), Internet: [www.gollaschconsulting.de](http://www.gollaschconsulting.de)*

<sup>2</sup>*AeT umweltplanung, Bismarckstraße 19, 56068 Koblenz Germany*

*E-mail: [nehring@aet-umweltplanung.de](mailto:nehring@aet-umweltplanung.de), Internet: [www.aet-umweltplanung.de](http://www.aet-umweltplanung.de)*

Received 5 November 2006; accepted in revised form 4 December 2006

### Abstract

More than 140 aquatic alien species (AAS) have been reported from coastlines of the North Sea and the Baltic Sea and from inland waters within the national borders of Germany. The majority of these species has established self-sustaining populations. The most important vectors of introduction are shipping, species imports for aquaculture purposes and species imports as part of the ornamental trade. Several AAS have reached German waters via shipping canals. Many species show a locally limited distribution, but almost half of all AAS have spread successfully across larger areas. Several introduced species are abundant and approximately 20 % of all AAS in Germany can be considered as invasive. Prime source regions are the north-western Atlantic, the Indo-Pacific, and the Ponto-Caspian region. For all source regions considered, the invasion rate has been increasing since the end of the last century.

*Key words:* Germany, North Sea, Baltic Sea, inland waters, aquatic species introductions, shipping, aquaculture, population status, invasive

### Introduction

Invasive alien species may threaten native species, alter habitats, and even affect ecosystem function (e.g. Eno et al. 1997, Nehring and Leuchs 1999, Wolff 2005), and thus represent a significant risk to the receiving environments. Following direct habitat destruction, invasive alien species are considered as the second most important cause of global biodiversity change (CBD 2000).

One of the first summaries of aquatic invaders in German coastal waters was prepared by Gollasch (1996). In 1997 Eno et al. published a summary of coastal aquatic alien species in the United Kingdom. Nehring and Leuchs (1999), Nehring (2000a), and Tittizer et al. (2000) published overviews on “neozoans” of the

German macroinvertebrate fauna. In 1999, Reise et al. published a summary of invasive species in the North Sea and several regional updates were published thereafter: e.g. Weidema (2000) for Nordic countries, Nehring (2005) and AeT umweltplanung (2006) for Germany, Jensen and Knudsen (2005) for Denmark and Wolff (2005) for The Netherlands. In 2006 Gollasch published an overview on introduced aquatic species known from European coasts. Here we update the earlier summaries of alien species in German inland and coastal waters. Another data set of aquatic invaders, which contains more comprehensive information for each species listed, is prepared by the authors for the currently ongoing EU-Programme Delivering Alien Invasive Species Inventories for Europe (DAISIE, see <http://www.daisie.se> for details).

### Aquatic alien species (AAS) in Germany


A total of 141 non-native taxa were reported from the waters considered in this overview, i.e. the coasts of the North Sea and the Baltic Sea and the inland waters within the national borders of Germany (Annex). The vast majority of these species were introduced by ship traffic and, intentionally, by stocking or for aquaculture. Species which reached the region on their own i.e. via drift with currents, swimming, or other ways of natural range expansion, were excluded from this overview. Most AAS have been reported from inland waters, followed by the coastal waters of the North Sea and the Baltic Sea.

More than two thirds of the known introduced species have established self-sustaining populations (Table 1). Some species were only recorded over a certain time period (e.g. the Hydrozoa *Bougainvillia macloviana* Lesson, 1830 the Anthozoa *Haliplanella luciae* (Verrill, 1898) and the Bivalvia *Crassostrea virginica* (Gmelin, 1791) and have since become extinct (Annex).

**Table 1.** Number of aquatic alien species (AAS) known from German waters. The number of AAS which are considered as established is listed separately

Region	All species	Established species
Baltic Sea	34	28
North Sea	62	49
Inland waters	86	82

Intentional fish introductions were predominantly motivated by a perceived improvement to the inland fisheries. About 70 “alien” fish species have been recorded in German waters (Geiter et al. 2002). A two century history of fish stocking and translocation makes it impossible to reconstruct the native range of most alien commercial fish species and their phylogeographic structure might also have been obscured. Consequently, some of these 70 species are considered as cryptogenic (see below). At present, ten fish species have been recognized as aliens and are established in self-sustaining populations with locally restricted distribution.



**Figure 1.** Number of aquatic species introductions into German waters for 20 year intervals between 1850 and 2006

The invasion rate has been increasing in all waters since the end of the last century, with the highest rate of increase found in inland waters (Figure 1). It is anticipated that more species were found in recent years as new findings are usually published with a time-lag. Many of the alien species are at least locally abundant (Table 2) and nearly half of all AAS have spread successfully across a larger area. A few alien species have developed large populations and mass developments have been observed as, for example, for the Chinese mitten crab *Eriocheir sinensis* Milne-Edwards, 1854 in German inland waters (Figure 2).

Major natural hydrographical and topographical differences exist between the three aquatic ecosystems considered (i.e., inland waters, North Sea and Baltic Sea coasts). These differences are also reflected in a distinct occurrence of alien species.

Some species were only found in single or a few records, i.e. the decapod *Callinectes sapidus* Rathbun, 1896, the anthozoan *Cereus pedunculatus* (Pennant, 1777), the hydroid *Gonionemus vertens* Agassiz, 1862, the horseshoe crab *Limulus polyphemus* Linnaeus, 1758 and the fish *Neogobius kessleri* (Günther, 1861) (Annex).

**Table 2.** Population status of aquatic alien species known from German waters

Population Status	Baltic Sea	North Sea	Inland Waters
Unknown	1	1	
Extinct	1	4	1
Single record(s)	1	4	
Rare	6	16	33
Local	11	16	29
Common	1	5	10
Abundant	13	16	14
Total	33	62	86

The relationship between salinity and species diversity is well known. In contrast to freshwater and pure seawater, brackish waters are characterized by the lowest number of indigenous species (“Artenminimum” sensu Remane 1934) and seem to provide opportunities for alien species invasions. It has recently been shown for German waters that the brackish areas of estuaries have been invaded more frequently by alien macroinvertebrate species than rocky or sandy sea shores or inland waters (Nehring 2006a). The author also stated that a low indigenous species richness in aquatic communities facilitates invasions of ‘new’ species, but the frequency and intensity (or size) of the inoculation are critical components in determining colonization success. Brackish waters seem to have many open ecological niches and are often exposed to intensive international ship traffic. Thus, these habitats have the highest potential for species introductions. In addition, estuaries are subjected to a two-sided invasion pressure by alien species via the ocean (e.g., due to shipping) and via inland waters (e.g., canal constructions).

Since the 1980s, polychaetes of the genus *Marenzelleria* have appeared in the North Sea and Baltic Sea. The taxonomic identification of the species, which were introduced with ballast water discharge of ocean going ships, was rather difficult and led to confusions and misidentifications. Sikorski and Bick (2004) showed that at least two *Marenzelleria* species occur in German waters: *Marenzelleria neglecta* Sikorski et Bick, 2004 and *M. viridis* (Verrill, 1873) (former taxonomic determinations and synonyms for *M. neglecta* are: *M. viridis*, *M. cf. viridis* and *M. Type II*, and for *M. viridis*: *M. wireni* Augener, 1913, *M. cf. wireni* and *M. Type I*). After the first appearance of *M. viridis* in 1979 in a Scottish estuary, the species arrived at the German North Sea coast in 1983 (Essink and Kleef 1986). First individuals of *M. viridis* were found in the German Baltic Sea by 2004 (Bastrop and Blank 2006). Since 1985 *M. neglecta* occurs along the German Baltic Sea coast (Bick and Burckhardt 1989). In 1996 the polychaete was detected in the Kiel Canal (connecting the North Sea with the Baltic Sea) as well as in the Elbe estuary (North Sea) and by 1997 *M. neglecta* had arrived in the Weser estuary (Nehring and Leuchs 2000). Both *Marenzelleria* species spread rapidly and became the predominant polychaete worms in German coastal waters. In the estuaries of the German North Sea coast both *Marenzelleria* species show distinct occurrences due to the salinity gradient (Nehring and Leuchs 2001). *M. viridis* prevails in the mesohaline zone, while *M. neglecta* colonizes mainly the oligohaline zone. In the area between the two zones, both species occur sympatrically.

While many alien species seem to remain insignificant additions to the native biota of Germany, approximately 20 % of the introduced species show invasive behaviour (sensu CBD 2000). The following provides a simple classification, modified after Jansson (1994) and Hopkins (2000), to document the different impacts of introduced alien species in the recipient ecosystem, viz:

- Disruption of existing interactions between species or food web links (e.g., predators, prey, grazers, and competition) - e.g. *Crassostrea gigas* (Thunberg, 1793) (Bivalvia), *Dikerogammarus villosus* (Sovinsky, 1894) (Amphipoda), *Dreissena polymorpha* (Pallas, 1771) (Bivalvia);
- Hybridisation with native and other alien species, resulting in changes of biological and


genetic diversity. Potential candidates in German waters: *Acipenser* spp. (Pisces), *Crassostrea gigas* (Bivalvia), *Lepomis* spp. (Pisces), *Spartina anglica* Hubbard, 1968 (Poacea);

- Introduction of parasites and disease agents. The introduced species may function as a host for pathogens or parasites which affect indigenous species - e.g., *Anguillicola crassus* (Kuwahara, Niimi et Hagaki, 1974) (Nematoda), *Orconectes limosus* (Rafinesque, 1817) (Decapoda);
- Habitat modification - e.g., *Chelicorophium curvispinum* (Sars, 1895) (Amphipoda), *Crassostrea gigas* (Bivalvia), *Hypania invalida* (Grube, 1860) (Polychaeta), *Sargassum muticum* (Yendo) Fensholt, 1955 (Phaeophyceae);
- Impact on species used in fisheries and aquaculture, resulting in decrease of output - e.g. *Anguillicola crassus* (Nematoda), *Eriocheir sinensis* (Decapoda), *Crassostrea gigas* (Bivalvia);
- Impact on resource users may result in harmful consequences on human health and well-being, recreation, and socio-economics - e.g., *Crassostrea gigas* (Bivalvia), *Elodea canadensis* Michaux, 1803 (Hydrocharitaceae), *Spartina anglica* (Poacea).



**Figure 2.** Mass upstream migration of juvenile Chinese mitten crabs, *Eriocheir sinensis*, in the Elbe River near Hamburg in 1998. Photo Stephan Gollasch

Although we have some information about some of the direct impacts of AAS, the longer-term ecological consequences for native plant and animal communities and the scale on which biodiversity is modified by invasive species is still poorly understood. Analyses of the economic effects of AAS are also needed.



**Figure 3.** Introduction vectors of aquatic alien species in German waters


The most important vectors for species introductions in the Baltic Sea and the North Sea are shipping and species imports for aquaculture. In inland waters most AAS invasions are attributed to canal constructions facilitating species migrations, to the release of species that have been imported with the ornamental trade, stocking and ship traffic (Figure 3).

Prime source regions for AAS that have invaded German waters are the Ponto-Caspian area, the north-western Atlantic and the Indo-Pacific for the Baltic Sea, the northern Pacific, the Indo-Pacific, the north-western Atlantic for the North Sea, and north America and the Ponto-Caspian area for inland waters (Figure 4).

#### Cryptogenic species

The native range of some of the species which have been considered as alien is controversial. These species are referred to as cryptogenic

species, i.e., species that are neither native or introduced (Carlton 1996). Those species include the polychaetes *Aphelochaeta marioni* (Saint-Joseph, 1894), *Microphthalmus similes* Bobretzky, 1870, *Nereis virens* Sars, 1835, *Polydora ligerica* (Ferronière, 1898) and the Dinophyceae *Prorocentrum redfieldii* Bursa, 1959 (Annex). As those species may be introduced, they were included in the Annex for reasons of comparison.



**Figure 4.** Source regions of aquatic alien species in German waters

Another interesting case is a turbellarian sampled from a ship hull in a German port. After careful taxonomic consideration it was found that this species is new to science and it was described as *Cryptostylochus hullensis* Faubel et Gollasch, 1996 (Polycladida, Acotylea, Plathelminthes). Because this flatworm is only known from this single sample, the native range remains unclear (Faubel and Gollasch 1996). The species was never found again in German waters

and is therefore not included in the species list attached.

#### *Species introductions and climate change*

We excluded species that reached German waters from their known distribution range by natural means such as range expansion from e.g., the north-east Atlantic or the Mediterranean Sea. It has been hypothesized that temporary or permanent climate change facilitates natural range expansion (Nehring 1998a, Stachowicz et al. 2002). Franke et al. (1999) and Franke and Gutow (2004) reported several nonindigenous species from the North Sea near Helgoland which are known to occur west of the British Channel and/or in the Mediterranean Sea. Examples include the decapods *Palaemon longirostris* Milne Edwards, 1837, *Portumnus latipes* (Pennant, 1777), the Polychaeta *Sabellaria alveolata* (Linnaeus, 1767) and the Bacillariophyceae *Thalassiosira hendeyi* Hasle et Fryxell, 1977.

The cord-grass *Spartina anglica*, a fertile hybrid of the European species *S. maritima* (Curtis) Fernald, 1916 and the North-American species *S. alterniflora* Loiseleur-Deslongchamps, 1807, was introduced into the Wadden Sea in the 1920s to promote sediment accumulation. However, the intended stabilization of mudflats was not always achieved. Recently this alien species has spread naturally into the tidal zone, where it displaces the native glass-word *Salicornia stricta* Dumort, 1868 (Figure 5). This range extension may have been promoted by higher spring temperatures. *S. anglica* may further benefit from climate change and may become more abundant in the near future, resulting in unforeseeable consequences for coastal protection (Nehring 2003, Nehring and Adersen 2006).

#### *Warm water effluents as hot spots of species invasions*

Alien species, native to warmer climate regimes, may also have colonised the North Sea in localities with unusual high water temperatures, e.g. near cooling water outlets of power plants. One example is the Pacific polychaete *Ficopomatus enigmaticus* (Fauvel, 1923). This brackish water species was first recorded in the London Docks, United Kingdom in 1922 (Eno et al. 1997), in the port of Vlissingen, The Netherlands in 1967 near a power plant (Wolff 2005) and also in the German Port of Emden in



close vicinity of a power plant (Kühl 1977a). Today, *F. enigmaticus* is widespread in coastal areas of all North Sea countries.

Another species which "benefited" from locally heated waters is the freshwater Asiatic clam *Corbicula fluminea* (O.F. Müller, 1756) which was first found in Europe in 1989 near the



**Figure 5.** The cord-grass *Spartina anglica* (in the background) displaces the native glass-wort *Salicornia stricta* (in the foreground), Wadden Sea near Eider estuary in 2004. Photo Stefan Nehring



**Figure 6.** Mass occurrence of the Asian clam *Corbicula fluminea* in the Rhine River near Koblenz in 2006. Photo Stefan Nehring

port of Rotterdam, The Netherlands (Wolff 2005). In 1990 it was collected from the German section of the Rhine River (Figure 6), in 1997 from the Danube, and in 1998 from the Elbe (Tittizer et al. 2000). It has been suggested that the successful dispersal of the Asiatic clam in European waters is correlated with winter water temperature minima of 2 °C (Schöll 2000). In Germany, temperatures of inland waters frequently drop below 2 °C in winter and consequently *C. fluminea* should have limited

opportunities for establishment. However, industrial and residential discharges of warm water into many German rivers have raised winter temperature almost permanently above 2°C, thereby promoting the establishment of *C. fluminea* in high abundances (Galil et al. 2007).

#### *Species findings attributed to drift*

Newly recorded species may also have reached German coastal regions by drift with exceptional water inflow due to rare hydrodynamic situations or storms. In some cases, such as for the Bacillariophyceae *Corethron criophilum* Castracane, 1886 and *Rhizosolenia indica* Peragallo, 1892, the Cirripedia *Lepas anatifera* Linnaeus, 1758 and *Lepas fascicularis* Ellis et Solander, 1786, the Decapoda *Pachygrapsus marmoratus* (Fabricius, 1787) and the clupeid fish *Sardina pilchardus* Walbaum, 1792 this has resulted in a temporary occurrence outside of their native range (Luther 1987, Nehring 1998b, Ehrich and Stransky 2001, G. Meurs (Nationalpark-Zentrum Multimar Wattforum, Tönning, Germany) pers. comm.). These species have not been included in the Annex.

In October 2006 the Ctenophore *Mnemiopsis leidyi* Agassiz, 1865 has been found for the first time along the German part of the Baltic Sea coast (U. Sommer and J. Javidpour (Leibniz-Institut für Meereswissenschaften, Kiel, Germany) pers. comm.). Recently it was also found in Dutch estuaries (Faasse and Bayha 2006, this issue), in the Skagerrak and Kattegatt (Hansson 2006, this issue) and in southern Norway (A. Jelmert (Floedevigen Research Station, His, Norway) pers. comm.), but not yet along the German North Sea coast. This western Atlantic species was possibly transported into the Baltic by easterly directed water currents or introduced by human activities, however, its current alien status is unknown.

#### *Canals as invasion corridors*

The natural barriers between river and sea basins that have existed since the end of the Pleistocene have been largely eliminated by canals built during the last centuries. The occurrence of 26 alien species in German waters can be attributed to canal construction. The following examples highlight the importance of shipping canals as invasion corridors.

The opening of the Bug-Prypjat Canal in 1784, which connects the Dnieper-Pripyat

system to the rivers Bug and Vistula, was of crucial importance for the early and frequent occurrence of Ponto-Caspian species in northern Europe (e.g., the invasive zebra mussel *Dreissena polymorpha*). After the opening of the Main-Danube Canal in Germany in 1992, which connects the Rhine River and the Danube River, this southern corridor is today the most important link between the Ponto-Caspian area and western Europe. Recently, several Ponto-Caspian species have been found in increasing abundances in the German rivers Main and Rhine (e.g. the isopod *Jaera istri* [Schleuter and Schleuter 1995]). In contrast Bernauer and Jansen (2006) reported that the polychaete *Hypania invalida* decreased in numbers in the upper Rhine River between 2003 and 2004.

In 1995 the Ponto-Caspian amphipod *Dikerogammarus villosus* arrived in the Rhine basin via the Main-Danube-Canal (Tittizer et al. 2000). Since then this new invader has dispersed over large distances in a short period of time and in 2000 the first organisms were observed in the German/Polish river Odra (Müller et al. 2001). This dynamic geographic expansion of *D. villosus* in Germany was facilitated by several man-made canals in northern Germany which created connections to all large river systems (Rhine, Weser, Elbe, Odra). Due to the rapidly increasing population density of this invasive amphipod it became a major component of the macrobenthic fauna in German freshwater systems, eliminating both native and other alien amphipod species (Tittizer et al. 2000, Haas et al. 2002, Nehring 2005).

More Ponto-Caspian species, mainly invertebrates and fishes, are expected to migrate into the North Sea basin via the Main-Danube-Canal. Especially those species which already occur in the upper and middle Danube will likely arrive in the North Sea basin soon.

The Chinese mitten crab (*Eriocheir sinensis*), introduced with ships and first recorded in the Aller River in 1912 (Schnakenbeck 1924, Marquard 1926), was reported from the North Sea coast in 1915 (Schnakenbeck 1924), and from the Baltic Sea in 1932 (Boettger 1933a, Peters 1933). *E. sinensis* was also found in the Kiel Canal in the 1920s (Neubaur 1926) and it is likely that the crab used the canal as the main invasion corridor to migrate from the North Sea into the Baltic Sea (Gollasch et al. 2006). Today the mitten crab can be found in the northern and easternmost parts of the Baltic (Ojaveer et al.

accepted). Using the same invasion corridor as *E. sinensis*, the north-American amphipod *Gammarus tigrinus* Sexton, 1939 may have reached the Baltic Sea from inland waters in the catchment of the North Sea, where it was intentionally introduced in 1957 (Schmitz 1960) and first recorded in Germany. *G. tigrinus* successfully spread and reached the North Sea coast by 1965 (Klein 1969) and the Baltic Sea in 1975 (Bulnheim 1976, 1980, Wawrzyniak-Wydrowska and Gruszka 2005).

An east-west (i.e., opposite to the more common direction of invasions) directed migration through the Kiel Canal may have occurred in the case of the decapod *Rhithropanopeus harrisi* (Gould, 1841). This crab was first recorded along the Baltic shores near Kiel and adjacent inland waters (Flehmuder Lake and Kiel Canal) in 1936 (Neubaur 1936), and it was subsequently recorded from the Wadden Sea (Cole 1982, Kühl 1977b, Adema 1991, Nehring 2000b, Van der Velde et al. 2000).

### Invasion Myths

The following section focuses on the importance of ships as species invasion vectors. Fact is that shipping continues since centuries and that ballast water is in use for more than 100 years. Some issues, formerly entitled "Invasions Myths" (J.T. Carlton (Williams College, Mystic, USA) pers. comm.), may arise:

*"All species, which could have been introduced, are here by now!"*

This is not the case. The "window of introduction" theory explains that all factors need to be right to enable a successful species introduction. These factors include e.g. temperature, salinity, food availability, lack of predators and the number of specimens for a founder population. It is believed that a successful invasion only occurs when all factors involved form the right environment for the candidate invader. However, the factors listed are highly varying and one can easily think of thousands of theoretical combinations indicating how rarely optimum conditions may occur in the receiving environment. Further, ship improvements result in larger ballast water tanks, more frequent ship arrivals and shorter voyage durations thereby increasing the survival rate of

species in transit. It should be noted that the zebra mussel was first recorded in the North American Great Lakes in the 1980s, but ships from its donor region arrived in the lakes since many decades before the species was introduced, i.e. it took quite some time and probably multiple discharge events until all factors triggering the invasion were right.

*“Why do we need to go active right now?”*

The number of invaders was increasing towards the end of the last century. Several investigations have shown that since 1950s the number of new records of invaders have clearly increased (Figure 1). Further new free trade agreements and ship improvements (see above) may have increased the invasion rate even further, thereby indicating the need for immediate action with the aim to reduce the number of new alien species arrivals.

*“Biological invasions are a natural phenomenon and happen anyway. The only thing we do is to speed up the process”*

This is simply not true as there is no natural means to transport a species from e.g. North America to Australia. Biogeographical textbooks describe the Pacific Ocean as a migration barrier as the duration of the zooplankton larval phase is too short to enable a distribution across the Pacific with natural means. Human mediated vectors, such as ballast water or hull fouling transports, are essential for a species to become dispersed across the Pacific. Also, freshwater species cannot reach new habitats separated by marine waters. However, ballast water releases from e.g. the freshwater port of St. Petersburg (Russia) in the port of Hamburg (Germany) may introduce species which could not reach the area by natural means due to the higher salinity in the western Baltic and North Sea.

*“Humans should not interfere with species distributions”*

Invasion biologists know that biological components and their interaction in an environment are not a stable process. It was also agreed that initiatives should not be undertaken to hinder natural migration activities of species. However, human mediated species introductions should be kept to a minimum as a precautionary approach. Case histories have shown severe,

unwanted impacts of invaders which were introduced unintentionally with e.g. ballast water or associated with aquaculture imports. Natural migrations and human mediated species introductions should clearly be treated separately.

*“Only 10% of the invaders show a significant impact”*

This statement refers to the "10s-rule". The rule was originally postulated based on invasion histories in terrestrial habitats. The figure was revised frequently. No matter how detailed these revisions were it has to be noted that each invasion has its impact on the recipient region. In some cases the impact is quite clear, in other instances the impact is not as obvious. Further, in many cases an impact is only noted when the invader forms a mass development which may occur long time after the initial introduction. In invasion biology it is not the quantity which matters, but the quality, i.e. just one introduced species may severely impact the receiving environment.

*“Phytoplankton species are not matter of discussion as these species are distributed world-wide anyway”*

It was documented that the number of phytoplankton blooms increased during the last two decades world-wide and it was suggested that this was supported not only by eutrophication but also by biological invasions. The recent occurrence of potentially toxin producing phytoplankton species in the North Sea is a good indication that we should be prepared for additional invaders of this kind.

*“Keep the ballast water onboard as long as possible and the species will die over time”*

Although many species die during the first days in ballast tanks, scientific studies have shown that after more than 4 months living zooplankton can be found in ballast tanks and under certain circumstances zooplankton species even reproduce in ballast tanks (Gollasch et al. 2000a,b, Gollasch et al. 2002). Further, some plankton species are enabled to form resting stages that survive unfavourable conditions for years or decades. Therefore, keeping ballast water onboard for longer periods of time is not a measure to significantly reduce the risk of species invasions.



“The exchange of ballast water in high seas is an appropriate means to reduce the number of invaders”

The exchange of ballast water in mid-ocean can reduce the abundance and diversity of taxa in ballast water. It is further unlikely that coastal organisms taken up in ports survive open ocean conditions where ballast water is exchanged – and plankton from high seas is unlikely to survive in coastal areas. In contrast to this assumption scientists showed that the exchange of ballast water could increase the species diversity in ballast tanks, especially in many domestic shipping routes, where no deep water exchange zones occur. Also the number of individuals in ballast tanks may increase when ballast water exchange is undertaken in zones with e.g. phytoplankton blooms. Ballast water exchange is therefore recommended as a very first management option, but effective treatment measures are urgently needed to avoid ballast water mediated species invasions in the future.

#### The future of alien species introductions into German waters and their potential impacts

The publication of recently introduced species in scientific journals is sometimes a time consuming process, and it is likely that by the time this checklist is published new alien species have already invaded German waters.

These may include the *Rapana venosa* (Valenciennes, 1846) (Gastropoda), which was observed for the first time in the south-western North Sea in 2005 (Kerckhof et al. 2006), but outside German national waters. This species was already known from European waters and the new findings in the North Sea likely represent a secondary introduction. However, the occurrence of this species in various locations in Europe may also be a result of multiple introductions from its native range. Noting its potential to spread, it is anticipated that this species may be found in German waters soon.

In 1999 Reise et al. prepared an overview of introduced species in the North Sea and concluded that most alien species can be considered "additive" and that they do not cause major unwanted impacts. However, there is evidence to the contrary. For example, the introduced Pacific oyster *Crassostrea gigas* is spreading in the Wadden Sea (Reise et al. 2005) with competitive effects on the *Mytilus edulis* Linnaeus, 1758 mussel beds (Figure 7). The

recent spread of *C. gigas* is likely triggered by (a) recent warm summers which support its recruitment and by (b) the absence of cold winters which promote recruitment of *M. edulis*. In northern Europe the Pacific oyster may benefit from global warming and may become more abundant than mussel beds have ever been (Diederich et al. 2005, Nehring 2006b).



Figure 7. Increasing abundance of the Pacific oyster *Crassostrea gigas* on a *Mytilus edulis* mussel bed near List, Sylt Island in 2005. Photo Stephan Gollasch

Because the impact of introduced species is potentially enormous, and very unpredictable, we should be aware of new species introductions. One known source of alien species is ship's ballast water which can contain millions of organisms and that is discharged in our coastal waters every day. Other vectors of introductions include species that are transported in the hull fouling of ships (Gollasch et al. 2000a, b, Nehring 2001, Gollasch 2002, Gollasch et al. 2002) and canal migrations. The latter have increased in magnitude and frequency over the past decade(s) (Harka and Biro 2004, Nehring 2005, Galil et al. 2007).

Similar to a worldwide trend, the rate of invasion of AAS has also increased in German waters since the 1950s, and will probably continue to rise due to the effects of climate

change and further improvements in ship design. It is hoped that measures, such as ballast water treatment to reduce the organism load or the installation of migration barriers such as deterrent electrical systems, salt or freshwater water locks, and air bubble curtains to reduce the uncontrolled range expansion of alien species via canals are taken soon to protect our waters from new species invasions and their potentially detrimental effects.

### Acknowledgements

The authors are grateful to a large number of colleagues who contributed to this inventory, but we cannot mention them individually here as the list would be too long. We also thank Dr. Wolfgang Jansen (North/South Consultants Inc., Winnipeg, Canada) for his critical review of the manuscript and for language editing. Gollasch is a contributing partner in the EU-funded DAISIE Programme (SSPI-CT-2003-511202, see also [www.daisie.se](http://www.daisie.se)).

### References

- Adema JPHM (1991) De krabben van Nederland en België (Crustacea, Decapoda, Brachyura). Nationaal Natuurhistorisch Museum, Leiden. 244 pp
- AeT umweltplanung (2006) Aquatic aliens in German inland and coastal waters. <http://www.aquatic-aliens.de>
- Ankel WE (1935) Die Pantoffelschnecke, ein Schädling der Auster. *Natur und Volk* 65: 173-176
- Arndt H and Schnese W (1986) Population dynamics and production of *Acartia tonsa* (Copepoda: Calanoida) in the Darss-Zingst estuary, southern Baltic. *Ophelia Suppl.* 4: 329-334
- Arndt W (1931) Die Tierwelt des Nordostseekanals und ihr Lebensraum. *Der Naturforscher* 8
- Arnold A (1990) Die neue Brehm-Bücherei. Eingebürgerte Fischarten. A. Ziemsen Verlag, Wittenberg Lutherstadt, 144 pp
- Ax P (1952) Eine Brackwasser-Lebensgemeinschaft an Holzpfehlern des Nord-Ostsee-Kanals. *Kieler Meeresforsch.* 9: 229-243
- Bartsch I and Kühlenkamp R (2000) The marine macroalgae of Helgoland (North Sea): an annotated list of records between 1845 and 1999. *Helgoland Marine Research* 54: 160-189
- Bastrop R and Blank M (2006) Multiple invasions - a polychaete genus enters the Baltic Sea. *Biol. Inv.* 18 (in press)
- Bastrop R, Röhner M, Sturmbauer C and Jürss K (1997) Where did *Marenzelleria* spp. (Polychaeta: Spionidae) in Europe come from? *Aquatic Ecol.* 31: 119-136
- Bäthe J (1995) Die Makroinvertebraten-Fauna der Weser. *Limnologie aktuell* 6: 175-190
- Bentheim-Jutting T Van (1922) Zoet en Brackwatermollusken. *Flora en fauna Zuiderzee*, Den Helder, 391-410
- Bernauer D, Kappus B and Jansen W (1996) Neozoen in Kraftwerksproben und Begleituntersuchungen am nördlichen Oberrhein. In: Gebhardt H, Kinzelbach R and Schmidt-Fischer S (eds) *Gebietsfremde Tierarten. Auswirkungen auf einheimische Arten, Lebensgemeinschaften und Biotope*. Ecomed, Landsberg, 87-96
- Bernauer D and Jansen W (2006) Recent invasions of exotic macroinvertebrates and loss of native species in the upper Rhine River, Germany. *Aquatic Invasions* 1: 55-71
- Bick A and Burckhardt R (1989) Erstnachweis von *Marenzelleria viridis* (Polychaeta Spionidae) für den Ostseeraum, mit einem Bestimmungsschlüssel der Spioniden der Ostsee. *Mitt. Zool. Mus. Berlin.* 65(2): 237-247
- Bick A and Zettler ML (1997) On the identity and distribution of two species of *Marenzelleria* (Polychaeta, Spionidae) in Europe and North America. *Aquatic Ecol.* 31: 137-148
- Bishop MWH (1947) Establishment of an immigrant barnacle in British coastal waters. *Nature (Lond.)* 159: 501-502
- Boettger CR (1933a) Die Ausbreitung der Wollhandkrabbe in Europa. *Sitzungsber. Ges. naturforsch. Freunde, Berlin* 1933, 399-415
- Boettger CR (1933b) Über die Ausbreitung der Muschel *Congeria cochleata* Nyst in europäischen Gewässern und ihr Auftreten im Nordostseekanal. *Zool. Anz.* 101: 43-48
- Boettger CR (1934) Der nordamerikanische Flußkrebs *Cambarus affinis* SAY in Deutschland. *Sitzungsber. Ges. naturforsch. Freunde*, 149-157
- Bousfield EL (1958) Fresh-water amphipod crustaceans of glaciated North America. *Canad. Field-Natural.* 72(2): 55-118
- Broch H (1924) Cirripedia thoracica von Norwegen und dem norwegischen Nordmeer. Eine systematische und biologisch-tiergeographische Studie. *Vidensk. Skr., Kistinia (Mat.-nat. Kl.)* 17: 1-121
- Buchmann K, Mellergaard S and Køie M (1987) *Pseudodactylogyrus* infections in eel: a review. *Diseases of Aquatic Organisms* 3: 51-57
- Buitendijk AM and Holthuis LB (1949) Note on the Zuiderzee Crab, *Rithropanopeus harrisi* (Gould) subspecies *tridentatus* (Maitland). *Zool. Meddelingen Rijksmuseum Natuurl. Hist. Leiden* 30: 95-106
- Bulnheim H-P (1976) *Gammarus tigrinus*, ein neues Faunenelement der Ostseeförde Schlei. *Schr. Naturw. Ver. Schlesw.-Holst.*, 46: 79-84
- Bulnheim H-P (1980) Zum Vorkommen von *Gammarus tigrinus* im Nord-Ostsee-Kanal. *Arch. Fischereiwiss.*, 30(1): 67-73
- Carlton JT (1996) Marine bioinvasions: The alteration of marine ecosystems by nonindigenous species. *Oceanography* 9: 1-7
- Caspers H (1950) Die Lebensgemeinschaft der Helgoländer Austernbank. *Helgoländer wiss. Meeresunters.* 3: 119-169
- CBD 2000. Global strategy on invasive alien species. Convention on Biological Diversity, UNEP/CBD/SBSTTA/6/INF/9, 1-52

Checklist for aquatic alien species in Germany

- Christiansen ME (1969) Decapoda Brachyura. Marine invertebrates of Scandinavia 2: 1-143 Universitetsforlaget, Oslo
- Cohen AN and Carlton JT (1995) Nonindigenous aquatic species in a United States estuary: a case study of the biological invasions of the San Francisco Bay and Delta. A Report for the US Fish and Wildlife Service, Washington D.C., and the National Sea Grant College Program, Connecticut Sea Grant, 201 pp
- Cole HA (1982) Status (1980) of Introductions of Non-indigenous Marine Species to North Atlantic Waters, ICES Cooperative Research Report 116, 87 pp
- Dahl F (1891) Untersuchungen über die Thierwelt der Unterelbe. Ber. Komm. wiss. Unters. deutsch. Meere Kiel 1887-1891. 6: 150-185
- De Lattin G (1967) Grundriß der Zoogeographie. Stuttgart, 386 pp
- Dehus P (1990) Die Verbreitung der Flußkrebse (Decapoda: Astacidae, Cambaridae) in Schleswig-Holstein. Faun.-Ökol. Mitt. 6: 95-105
- Diederich S, Nehls G, van Beusekom JEE and Reise K (2005) Introduced Pacific oysters (*Crassostrea gigas*) in the northern Wadden Sea: invasion accelerated by warm summers? Helgol. Mar. Res. 59: 97-106
- Dijkema KA (1983) The salt-marsh vegetation of the mainland coast, estuaries and Halligen. In: Dijkema KA and Wolff WJ (eds) Flora and vegetation of the Wadden Sea islands and coastal areas. Stichting, Leiden. 185-220
- Duncker G (1939) Hundsfische (*Umbra spec.*) in Schleswig Holstein? Heimat (Kiel) 49: 300-301
- Eggers TO and Anlauf A (2005) *Obesogammarus crassus* (G.O. Sars, 1894) (Crustacea: Amphipoda) erreicht die Elbe. Lauterbornia 55: 125-128
- Eggers TO and Martens A (2004) Bestimmungsschlüssel der Süßwasser-Amphipoda (Crustacea) Deutschlands. Lauterbornia 42: 1-68
- Ehrich S and Stransky (2001) Spatial and temporal changes in the southern species component of North Sea bottom fish assemblages. Senckenbergiana marit. 31: 143-150
- Elbrächter M (1994) Phytoplankton und toxische Algen im Wattenmeer. In: Lozán JL, Rachor E, Reise K, Westenhagen Hv and Lenz W (eds), Warnsignale aus dem Wattenmeer. Blackwell Berlin: 81-86
- Elbrächter M (1999) Exotic flagellates of coastal North Sea waters. Helgoländer Meeresunters. 52: 235-242
- Eno NC, Clark RA and Sanderson WG (eds.) (1997) Non-native Species in British Waters: a Review and Directory. Joint Nature Conservation Committee, Peterborough, 152 pp
- Essink K and Kleef HL (1986) Establishment of a population of the Spionid worm *Marenzelleria* in the Ems Estuary (the Netherlands, Fed. Rep. of Germany). Proceedings of COST-647 Workshop "Intertidal Soft Sediments", Dec. 1986, St. Valery-sur-Somme, France. 5 pp
- Essink K (1985) On the appearance of the American jack-knife clam *Ensis directus* (Conrad, 1883) (Mollusca, Cultellidae) in the Dutch Wadden Sea. Basteria 49(4-6): 73-80
- Faasse M (1998) Vindplaatsen van het mosdiertje *Bugula stolonifera* Ryland, 1960 in Nederland. Zeepaard 58: 48-51
- Faasse MA and Bayha KM (2006) The ctenophore *Mnemiopsis leidyi* A. Agassiz 1865 in coastal waters of the Netherlands: an unrecognized invasion? Aquatic Invasions 4: 270-277
- Faubel A and Gollasch S (1996) *Cryptostylochus hullensis* n. sp. (Polycladida, Acotylea, Plathelminthes): a possible case of transoceanic dispersal on a ships hull. Helgoländer Meeresunters. 50: 533-537
- Fletcher RL and Farrell P (1999) Introduced brown algae in the North East Atlantic, with particular respect to *Undaria pinnatifida* (Harvey) Suringar. Helgoländer Meeresunters. 52: 259-275
- Franke H-D and Gutow L (2004) Long-term changes in the macrozoobenthos around the rocky island of Helgoland (German Bight, North Sea). Helgol. Mar. Res. 58: 303-310
- Franke HD, Gutow L and Janke M (1999) The recent arrival of the oceanic isopod *Idotea metallica* Bosc off Helgoland (German Bight, North Sea): an indication of a warming trend in the North Sea? Helgoländer Meeresunters. 52: 347-357
- Franz HW (1992) Der Rhein und seine Besiedlung im Wandel: Schwebstoffzehrende Organismen (Hydrozoa, Kaptozoa und Bryozoa) als Indikatoren für den ökologischen Zustand eines Gewässers. Pollichia-Buch 25: 1-167
- Galil BS, Nehring S and Panov V (2007) Waterways as invasion highways - Impacts of climate change and globalization. In: Nentwig W (ed.) Biological invasions. Ecological Studies Vol. 193, Springer, Berlin: 59-74
- Geissen HP and Schöll F (1998) Erste Nachweise des Fischegels *Caspiobdella fadejewi* (Epstein 1961) (Hirudinea: Piscicolidae) im Rhein. Lauterbornia 33: 11-12
- Geiter O (1998) Aktuelle Meldungen. Neozoen Newsletter 2: 10
- Geiter O, Homma S and Kinzelbach R (2002) Bestandsaufnahme und Bewertung von Neozoen in Deutschland. Umweltbundesamt Texte, 25/02: 173 pp (plus appendices)
- Gerstmeier R and Romig T (1998) Die Süßwasserfische Europas: für Naturfreunde und Angler. Franckh-Kosmos Verlag, Stuttgart, Germany. 368 pp
- Gessner J, Debus L, Filipiak J, Spratte S, Skora KE and Arndt G-M (1999) Development of sturgeon catches in German and adjacent waters since 1980. Journal of Applied Ichthyology 15: 136-141
- Gollasch S and Riemann-Zürneck K (1996) Transoceanic dispersal of benthic macrofauna: *Haliplanella luciae* (Verrill, 1898) (Anthozoa, Actinaria) found on a ship's hull in a shipyard dock in Hamburg Harbour, Germany. Helgoländer Meeresunters. 50: 253-258
- Gollasch S (1996) Untersuchungen des Arteintrages durch den internationalen Schiffsverkehr unter besonderer Berücksichtigung nichtheimischer Arten. Diss., Univ. Hamburg; Verlag Dr. Kovac, Hamburg, 314 pp
- Gollasch S, Lenz J, Dammer M and Andres HG (2000a) Survival of tropical ballast water organisms during a cruise from the Indian Ocean to the North Sea. J. Plankton Res. 22(5): 923-937
- Gollasch S, Rosenthal H, Botnen H, Hamer J, Laing I, Leppäkoski E, Macdonald E, Minchin D, Nauke M, Olenin S, Utting S, Voigt M and Wallentinus I (2000b) Fluctuations of zooplankton taxa in ballast water during

- short-term and long-term ocean-going voyages. *Internat. Rev. Hydrobiol.* 85(5-6): 597-608
- Gollasch S (2002) The importance of ship hull fouling as a vector of species introductions into the North Sea. *Biofouling* 18(2): 105-121
- Gollasch S, Macdonald E, Belson S, Botnen H, Christensen J, Hamer J, Houvenaghel G, Jelmert A, Lucas I, Masson D, McCollin T, Olenin S, Persson A, Wallentinus I, Wetsteyn B and Wittling T (2002) Life in Ballast Tanks. 217-231 pp. In: Leppäkoski E, Gollasch S and Olenin S (eds.): *Invasive Aquatic Species of Europe: Distribution, Impacts and Management*. KLUWER Academic Publishers, Dordrecht, The Netherlands. 583 pp
- Gollasch S (2006) Overview on Introduced Aquatic Species in European Navigational and Adjacent Waters. *Helgol. Mar. Res.* 60: 84-89
- Gollasch S, Galil BS and Cohen A (eds.) (2006). *Bridging Divides – Maritime Canals as Invasion Corridors*. SPRINGER Academic Publishers, Dordrecht, The Netherlands. 315 pp
- González-Ortegón E, Cuesta JA and Schubart CD (2006) First report of the oriental shrimp *Palaemon macrodactylus* Rathbun, 1902 (Decapoda, Caridea, Palaemonidae) from German waters. *Helgol. Mar. Res.*, online, visited 2006-09-12
- Gruner H-E (1962) Schwimmkrabben. In: Pax F (ed.) *Meeresprodukte*. Gebrüder Borntraeger, Berlin, 336-340
- Gruner H-E (1965) Krebstiere oder Crustacea. V: Isopoda. In: Dahl F (ed.). *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise*. Teil 51 und 53. Verlag G. Fischer, Jena, 380 pp
- Gruzka P (1999) The Odra River Estuary as a gateway for alien species immigration to the Baltic Sea basin. *Acta hydrochim. hydrobiol.* 27(5): 374-382
- Gugel J (1995) Erstnachweis von *Eunapius carteri* (Bowerbank, 1863) (Porifera: Spongillidae) für Mitteleuropa. *Lauterbornia* 20: 103-109
- Gürs I, Gürs K, Paustian J and Wiese V (1993) *Ensis directus* (Conrad 1843) zum ersten Mal in der südwestlichen Ostsee (Bivalvia: Solenidae). *Schriften aus dem Malakozoologischen Haus der Natur* 6: 79
- Haas G, Brunke M and Streit B (2002) Fast turnover in dominance of exotic species in the Rhine river determines biodiversity and ecosystem function: An affair between amphipods and mussels. In: Leppäkoski E, Gollasch S and Olenin S (eds.) *Invasive aquatic species of Europe - Distribution, impacts and management*. Kluwer, Dordrecht, 426-432
- Hagmeier A and Kändler R (1927) Neue Untersuchungen im Nordfriesischen Wattenmeer und auf den fiskalischen Austernbänken. *Wiss. Meeresunters. Abt. Helgoland* 16(6): 1-90
- Hahn A (1956) Die Bekämpfung der Bohrmuschel. *Die Küste* 5: 49-72
- Hansson HG (2006) Ctenophores of the Baltic and adjacent Seas – the invader *Mnemiopsis* is here! *Aquatic Invasions* 4: xx-xx
- Harka Á and Biro P (2004) Role of global warming and canalisation in the Central-European distribution of certain Ponto-Caspic fish species. p. 73 in *Abstracts of the XI European Congress of Ichthyology (ECI XI)*. Estonian Marine Institute Report Series No. 12
- Hartlaub C (1897) *Die Hydromedusen Helgolands*. *Wiss. Meeresunters. Abt. Helgoland*, II, 449-536
- Hartlaub C (1911) *Craspedote Medusen, Fam. III, Margelidae*. – In: Brandt K and Apstein C (eds.) *Nordisches Plankton*, Teil XII/1, Lief. 2. Lipsius and Tischer, Kiel Leipzig, 137–236
- Hartmann-Schröder G and Stripp K (1968) Beiträge zur Polychaetenfauna der Deutschen Bucht. *Veröffentl. Inst. Meeresforsch. Bremerhaven* 11: 1-24
- Hartmann-Schröder G (1996) Annelida, Borstenwürmer, Polychaeta. In: Dahl F (ed.) *Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise*, Teil 58, 2. Aufl. Verlag G. Fischer, Jena, 648 pp
- Hasle GR (1983) *Thalassiosira punctigera* (Castr.) comb. nov., a widely distributed marine planktonic diatom. *Nordic Journal of Botany* 3: 593-608
- Hasle GR (1990) Diatoms of the Oslo Fjord and Skagerrak. Species new to the area: immigrants or overlooked in the past? *Blyttia* 48: 33-38
- Hauer J (1950) Der nordamerikanische Strudelwurm *Euplanaria tigrina* (Girard) am Oberrhein. *Beiträge naturk. Forsch. SW.-Deutschlands* 9(1): 70-75
- Hauser B (1973) Bestandsänderungen der Makrofauna an einer Station im ostfriesischen Watt. *Jber. Forschungsstelle Norderney* 24: 171-203
- Havinga B (1929) *Krebse und Weichtiere*. In: Lübbert H and Ehrenbaum E (eds.), *Handbuch der Seefischerei Nordeuropas*, Band III, Heft 2. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 147 pp
- Hickel W, Hagmeier E and Drebes G (1971) *Gymnodinium* blooms in the Helgoland Bight (North Sea) during August 1968. *Helgoländer wiss. Meeresunters.* 22: 401-416
- Hinkelmann A (1899) Die Fische und sonstigen Nutztiere des Kaiser-Wilhelm-Kanals mit besonderer Berücksichtigung der Lebensverhältnisse des Herings. *Die Heimat*. 9: 173-178
- Hoek PPC (1875) *Eerste bijdrage tot een nauwkeuriger kennis der sessiele Cirripeden*. PhD thesis, University of Leiden. 94 pp
- Holthuis LB (1950) Decapoda. A. Natantia, Macrura Reptantia, Anomura en Stomatopoda. *Fauna van Nederland* 15: 1-120
- Hopkins CCE (2000) A review of introductions and transfers of alien marine species in the North Sea area. Report to the Norwegian Ministry of the Environment. *AquaMarine Advisers*, 74 pp
- Hopkins CCE (2001) Actual and potential effects of introduced marine organisms in Norwegian waters, including Svalbard. *Research Report 2001-1*, 53 pp. Norwegian Directorate for Nature Management.
- Huber MGJ and Schubart CD (2005) Distribution and reproductive biology of *Austropotamobius torrentium* in Bavaria and documentation of a contact zone with the alien crayfish *Pacifastacus leniusculus*. *Bull. Fr. Pêche Piscic.* 376/377: 759-776
- Hussner A (2005) *Aquatische Neophyten in NRW*. <http://www.aquatischeneophyten.de>

Checklist for aquatic alien species in Germany

- Jaeckel S (1962) Die Tierwelt der Schlei – Übersicht einer Brackwasserfauna. Schr. Naturwiss. Vereins Schleswig-Holstein 33: 11-32
- Jansson K (1994) Alien species in the marine environment: Introductions to the Baltic Sea and the Swedish west coast. Swedish Environmental Protection Agency Report No. 4357, 68 pp
- Jensen K and Knudsen J (2005) A summary of alien marine invertebrates in Danish waters. Oceanological and Hydrobiological Studies 34(Suppl 1): 137-162
- Jungbluth JH (1996) Einwanderer in der Molluskenfauna von Deutschland. I. Der chorologische Befund. In: Gebhardt H, Kinzelbach R and Schmidt-Fischer S (eds.) Gebietsfremde Tierarten. Ecomed, Landsberg, 105-125
- Kerckhof F, Vink RJ, Nieweg DC and Post JJJ (2006) The veined whelk *Rapana venosa* has reached the North Sea. Aquatic Invasions 1: 35-37
- Ketelaars HAM, Lambregts-van de Clundert FE, Carpentier CJ, Wagenvoort AJ and Hoogenboezem W (1999) Ecological effects of the mass occurrence of the Ponto-Caspian invader, *Hemimysis anomala* G.O. Sars, 1907 (Crustacea: Mysidacea), in a freshwater storage reservoir in the Netherlands, with notes on its autecology and new records. Hydrobiologia 394: 233-248
- Kinzelbach R (1991) Die Körbchenmuscheln *Corbicula fluminalis*, *Corbicula fluminea* und *Corbicula fluviatilis* in Europa (Bivalvia: Corbiculidae). Mainer Naturwiss. Arch. 29: 215-228
- Kirchenpauer JU (1862) Die Seetonnen der Elbmündung. Abhandl. Gebiete Naturwiss. 4(3): 1-59
- Klein G (1969) Amphipoden aus der Wesermündung und der Helgoländer Bucht, mit Beschreibung von *Talorchestia frisiae* n.sp. Veröff. Inst. Meeresforsch. Bremerhaven 11: 173-194
- Klie W (1933) Neues zur Crustaceen-Fauna Nordwestdeutschlands. Abh. Nat. Ver. Bremen, 28 (4): 271-276
- Kluijver MJ de (1991) Sublittoral hard substrate communities off Helgoland. Helgoländer Meeresunters. 45: 317-344
- Knudsen J (1989) Immigration of marine invertebrates to the Limfjord (Denmark) and the North Sea - Baltic transition area. In: Spanier E, Steinberger Y and Luria M (eds.) Environmental Quality and Ecosystem Stability. Vol. IV-B, Environmental Quality. ISEEQ Publication, Jerusalem, 135-145
- Kolumbe E (1931) *Spartina townsendii* - Anpflanzungen im schleswig-holsteinischen Wattenmeer. Wissenschaftliche Meeresuntersuchungen, Abt. Kiel, N.F. 21: 66-73
- Kothe P (1968) *Hypania invalida* (Polychaeta, Sedentaria) und *Jaera istri* (Isopoda) erstmals on der deutschen Donau. Arch. Hydrobiol. Suppl. 34: 88-114
- Kowarik I (2003) Biologische Invasionen - Neophyten und Neozoen in Mitteleuropa. Ulmer Stuttgart, 380 pp
- Kraeplin K (1887) The German freshwater Bryozoa. A monography. Abhandl. Verhandl. Naturwiss. Vereins Hamburg 10: 1-168
- Kuckuck P (1957) Der Strandwanderer. Lehmanns Verlag, München, 95 pp
- Kühl H (1954) Über das Auftreten von *Elminius modestus* Darwin in der Elbmündung. Helgoländer wiss. Meeresunters. 5: 53-56
- Kühl H (1962) Die Hydromedusen der Elbmündung. Abhandl. Verhandl. Naturwiss. Vereins in Hamburg. 6: 209-232
- Kühl H (1965) Fang einer Blaukrabbe, *Callinectes sapidus* Rathbun (Crustacea, Portunidae) in der Elbmündung. Arch. Fischereiwiss., 15(3): 225-227
- Kühl H (1977a) *Mercierella enigmatica* (Polychaeta: Serpulidae) an der deutschen Nordseeküste. Veröff. Inst. Meeresf. Bremerhaven 16: 99-104
- Kühl H (1977b) Neue Tierarten an der deutschen Nordseeküste. Inf. Fischw. 24(2): 61-62
- Kylin H (1930) Über die Entwicklungsgeschichte der Florideen. Lunds Univ. Årsskr. Swed. 26(6): 1-104
- Laing I (1999) *Coscinodiscus wailesii*. In: Gollasch S, Minchin D, Rosenthal H and Voigt M (eds) Exotics Across the Ocean. Case Histories on Introduced Species. Logos Verlag, Berlin
- Laufer H (2004) Zum Beutespektrum einer Population von Ochsenfröschen nördlich von Karlsruhe. Faunistische Abhandlungen 25: 139-150
- Lloyd WA (1874) On the occurrence of *Limulus polyphemus* off the coast of Holland, and on the transmission of aquarium animals. Zoologist, Series 2(9): 3845-3855
- Lu D and Göbel J (2000) *Chattonella* sp. bloom in North Sea, spring 2000. Harmful Algae News 21:10-11
- Luther G (1987) Seepocken der deutschen Küstengewässer. Helgoländer Meeresunters. 41: 1-43
- Maggs CA and Stegenga H (1999) Red algal exotics on North Sea coasts. Helgoländer Meeresuntersuch. 52: 243-258
- Marquard O (1926) Die Chinesische Wollhandkrabbe, *Eriocheir sinensis* Milne- Edwards, ein neuer Bewohner deutscher Flüsse. Fischerei 24: 417-433
- Martens A, Gerecke R and Grabow K (2006) *Caspihalacarus hyrcanus* (Acari) im Oberrhein. Der erste Fund einer neozoischen Wassermilbe in Deutschland und Frankreich. Lauterbornia 56: 27-34
- Martens A, Grabow K, Rapp S and Schoolmann G (in press) Der Krebsigel *Xironogiton victoriensis* wurde mit dem Signalkrebs *Pacifastacus leniusculus* auch nach Deutschland eingeschleppt (Annelida: Branchiobdellida; Crustacea: Astacidae). Lauterbornia
- Meister A (1997) Lebenszyklus, Autökologie und Populationsökologie der Körbchenmuscheln *Corbicula fluminea* und *Corbicula fluminalis* (Bivalvia, Corbiculidae) im Inselrhein. Umweltplanung, Arbeits- und Umweltschutz (Schriftenreihe Hessisches Landesamt für Umwelt, Wiesbaden) Heft 238, 170 pp
- Meyer-Waarden PE (1964) Mastversuche mit portugiesischen Austern im deutschen Wattengebiet. Inf. Fischw. 11: 77-78
- Millar RH (1960) The identity of the Ascidians *Styela manniculata* Carlisle and *S. clava* Herdman. J. Mar. Biol. Ass. U.K. 39: 509-511
- Minchin D and Rosenthal H (2002) Exotics for Stocking and Aquaculture, Making Correct Decisions. In: Leppäkoski E, Gollasch S and Olenin S (eds.) Invasive Aquatic Species of Europe: Distribution, Impacts and Management. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 206-216
- Minchin D, McGrath D and Duggan CB (1995) The Slipper Limpet, *Crepidula fornicata* (L.), in Irish waters, with a review of its occurrence in the North-Eastern Atlantic. J. Conch., London 35: 249-256



- Möbius K (1887) Schlußbericht über den Versuch des deutschen Fischereivereins, kanadische Austern in der Ostsee anzusiedeln. Deutscher Fischerei Verein. Mitt. Section Küsten- und Hochseefisch. 1
- Müllegger S (1921) *Heliactis bellis* bei Büsum gefunden. Schrift. Zool. Stat. Büsum, 2: 61
- Müller O, Zettler ML and Gruszka P (2001) Verbreitung und Status von *Dikerogammarus villosus* (Sovinski 1894) (Crustacea: Amphipoda) in der mittleren und unteren Strom-Oder und den angrenzenden Wasserstraßen. *Lauterbornia* 41: 105-112
- Muus BJ and Dahlström P (1968) Süßwasserfische. BLV Verlagsgesellschaft, München. 224 pp
- Nehls G (2004) *Gracilaria vermiculophylla* (Gracilariales, Rhodophyta) – a Pacific algae spreading on tidal flats of the Wadden Sea. Presentation at MARBEF Workshop: Aquatic invasive species and the functioning of European coastal ecosystems. Jan. 2005, List, Sylt (Germany)
- Nehring S (1998a) Establishment of thermophilic phytoplankton species in the North Sea - biological indicators of climatic changes? *ICES J. Mar. Sci.* 55: 818-823
- Nehring S (1998b) Non-indigenous phytoplankton species in the North Sea: supposed region of origin and possible transport vector. *Archive of Fishery and Marine Research* 46: 181-194
- Nehring S (2000a) Neozoen im Makrozoobenthos der deutschen Ostseeküste. *Lauterbornia* 39: 117-126
- Nehring S (2000b) Zur Bestandssituation von *Rhithropanopeus harrisi* (Gould, 1841) in deutschen Gewässern: die sukzessive Ausbreitung eines amerikanischen Neozoons (Crustacea: Decapoda: Panopeidae). *Senckenbergiana marit.* 30: 115-122
- Nehring S (2001) After the TBT era: Alternative anti-fouling paints and their ecological risks. *Senckenbergiana marit.* 31: 341-351
- Nehring S (2002) Biological invasions into German waters: an evaluation of the importance of different human-mediated vectors for nonindigenous macrozoobenthic species. In: Leppäkoski E, Gollasch S and Olenin S (eds.) *Invasive aquatic species of Europe - Distribution, impacts and management*. Kluwer, Dordrecht, 373-383
- Nehring S (2003) Alien species in the North Sea: invasion success and climate warming. *Ocean Challenge* 13 (3): 12-16
- Nehring S (2005) International shipping - A risk for aquatic biodiversity in Germany. In: Nentwig W, Bacher S, Cock MJW, Dietz H, Gigon A and Wittenberg R (eds.) *Biological Invasions - From Ecology to Control*. *Neobiota* 6: 125-143
- Nehring S (2006a) Four arguments why so many alien species settle into estuaries, with special reference to the German river Elbe. *Helgol. Mar. Res.* 60: 127-134
- Nehring S (2006b) NOBANIS - Invasive Alien Species Fact Sheet - *Crassostrea gigas*. Online Database of the North European and Baltic Network on Invasive Alien Species - NOBANIS [www.nobanis.org](http://www.nobanis.org), 14 pp
- Nehring S (2006c) The Ponto-Caspian amphipod *Obesogammarus obesus* (Sars, 1894) arrived the Rhine River via the Main-Danube Canal. *Aquatic Invasions* 1: 148-153
- Nehring S and Adersen H (2006) NOBANIS - Invasive Alien Species Fact Sheet - *Spartina anglica*. Online Database of the North European and Baltic Network on Invasive Alien Species - NOBANIS [www.nobanis.org](http://www.nobanis.org), 12 pp
- Nehring S and Leuchs H (1999) Neozoa (Makrozoobenthos) an der deutschen Nordseeküste - Eine Übersicht. Bundesanstalt für Gewässerkunde Koblenz, Bericht BfG-1200, 131 pp
- Nehring S and Leuchs H (2000) Neozoen im Makrozoobenthos der Brackgewässer an der deutschen Nordseeküste. *Lauterbornia* 39: 73-116
- Nehring S and Leuchs H (2001) Das BfG-Ästuarmonitoring Makrozoobenthos in Eider, Elbe, Ems, Jade und Weser. Jahresbericht 2000 der Bundesanstalt für Gewässerkunde, Koblenz: 93-95
- Neubaur R (1926) Eine chinesische Wollhandkrabbe im Wittensee. *Fischereizeitung*. (Neudamm). 29: 1111
- Neubaur R (1936) Ein neuer Mitbewohner schleswig-holsteinischer Fischgewässer. *Fisch.-zeit.* 39: 725-726
- Neuhaus E (1933) Studien über das Stettiner Haff und seine Nebengewässer. *Zeitschr. Fisch. u. deren Hilfswiss.* 31: 427-489
- Ojaveer H, Gollasch S, Jaanus A, Kotta J, Laine AO, Minde A, Normant M and Panov V (in press) Chinese mitten crab *Eriocheir sinensis* (H. Milne-Edwards, 1853) (Crustacea, Decapoda, Varunidae) population in the Baltic Sea – a supply-side invader? *Biological Invasions* (accepted)
- Ostenfeld CH (1908) On the immigration of *Biddulphia sinensis* Grev. and its occurrence in the North Sea during 1903-1907. *Medd. Komm. Havunders., Ser. Plankton*, 1(6): 1-46
- Panning A (1938) The Chinese Mitten Crab. *Smithsonian Rep.*, 361-375
- Pax F (1920) Die Aktinienfauna von Büsum. *Schr. Zool. Stat. Büsum Meeresk.*, 5: 1-24
- Pax F (1936) Anthozoa. In: Grimpe G and Wagler E (eds.) *Die Tierwelt der Nord- und Ostsee*. Bd. 30, Akad. Verlagsges. Becker and Erler, Leipzig, 317 pp
- Peters N (1933) B. Lebenskundlicher Teil. In: Peters N and Panning A (eds.) *Die chinesische Wollhandkrabbe (Eriocheir sinensis) H. MILNE-EDWARDS* in Deutschland. Akademische Verlagsgesellschaft mbH, Leipzig, 59-156
- Peters N and Hoppe H (1938) Über Bekämpfung und Verwertung der Wollhandkrabbe. *Mitt. Hamb. zool. Mus. Inst., Hamburg* 47: 140-171
- Petersen KS, Rasmussen KL, Heinemeier J and Rud N (1992) Clams before Columbus? *Nature* 359: 679
- Podraza P, Ehlert T and Roos P (2001) Erstnachweis von *Echinogammarus trichiatus* (Crustacea: Amphipoda) im Rhein. *Lauterbornia* 41: 129-133
- Post D and Landmann M (1994) Verbreitungsatlas der Fließgewässerfauna in Ostfriesland. *Staatl. Amt Wasser Abfall*, 141 pp
- Potel S, Geissen HP and Dohmen GP (1998) Erste Nachweise von *Barbronia weberi* (Blanchard 1897) (Hirudinea: Salifidae) im deutschen Rheingebiet. *Lauterbornia* 33: 1-4
- Rady GA (1913) Anleitung zu den Grundzügen für die Entwicklung einer Austernkultur an unserer deutschen Nordseeküste. Lehren aus der Zeit zur Hebung der Küstenfischerei. Hans Hofmann, Norden.



Checklist for aquatic alien species in Germany

- Reibisch J (1926) Über Änderungen in der Fauna der Kieler Bucht. Schr. naturw. Ver. Schleswig-Holstein 17: 227-232
- Reise K (1998a) Exoten der Nordseeküste. Wattenmeer Internat. 1: 21-22
- Reise K (1998b) Pacific Oysters invade mussel beds in the European Wadden Sea. Senckenbergiana marit., 28(4/6): 167-175
- Reise K, Dankers N, Essink K (2005) Introduced species. In: Essink, K. et al. (eds.) Wadden Sea Quality Status Report 2004. Wadden Sea Ecosystem No.19. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, 155-161
- Reise K, Gollasch S and Wolff WJ (1999) Introduced marine species of the North Sea coasts. Helgoländer Meeresunters. 52: 219-234
- Remane A (1934) Die Brackwasserfauna. Verh. dt. zool. Gesell. 36: 34-74
- Rudolph K (1997) Zum Vorkommen des Flohkrebse *Pontogammarus robustoides* im Peenemündungsgebiet. Natur und Museum 127: 306-312
- Schadt J (2000) Neue Fischart im Main entdeckt: Marmorierter Grundel (*Proterorhinos marmoratus*). Fischer & Teichwirt 6/2000: 217-218
- Scheibel W (1974) *Ameira divagans* Nicholls, 1939 (Copepoda Harpacticoida). Mikrofauna Meeresboden 38: 1-10
- Schellenberg A (1928) Krebstiere oder Crustacea. II: Decapoda, Zehnfüßer. In: Dahl F (ed.) Die Tierwelt Deutschlands und der angrenzenden Meeresteile nach ihren Merkmalen und nach ihrer Lebensweise, Teil 10. Verlag G. Fischer, Jena, 146 pp
- Schlesch H (1932) Über die Einwanderung nordamerikanischer Meeresmollusken in Europa unter Berücksichtigung von *Petricola pholadiformis* Lam. und ihrer Verbreitung im dänischen Gebiet. Arch. Moll. 64: 146-154
- Schimmer H (1995) Erstnachweis von *Piscicola haranti* (Jarry 1960) (Hirudinea) in Nordrhein-Westfalen. Lauterbornia 20: 111-113
- Schleuter M and Schleuter A (1995) *Jaera istri* VEUILLE (Janiridae, Isopoda) aus der Donau erreicht über den Main-Donau-Kanal den Main. Lauterbornia 21: 177-178
- Schlienz W (1922) Eine Süßwasser-*Orchestia* in der Außenalster in Hamburg. Arch. Hydrobiol. 14: 144-150
- Schmitz W (1960) Die Einbürgerung von *Gammarus tigrinus* Sexton auf dem europäischen Kontinent. Arch. Hydrobiol. 57: 223-225
- Schnakenbeck W (1924) Ueber das Auftreten chinesischer Krabben in der Unterelbe. Schr. für Süßwasser- und Meereskunde 2(5): 125-129
- Schöll F (2000) Die Wassertemperatur als verbreitungsregulierender Faktor von *Corbicula fluminea* (O.F. Müller 1774). Hydrol. Wasserbewirtschaft. 44: 318-321
- Schories D and Selig U (2006) Die Bedeutung eingeschleppter Arten (alien species) für die Europäische Wasserrahmenrichtlinie am Beispiel der Ostsee. Rostock. Meeresbiolog. Beitr. 15: 147-158
- Schrey I und Buschbaum C (2006) Asiatische Gespensterkrebse (*Caprella mutica*) erobern das deutsche Wattenmeer. Natur- und Umweltschutz (Zeitschrift Mellumrat) 5: 26-30
- Schulze FE (1981) Über den Bau und die Entwicklung von *Cordylophora lacustris* Allmann nebst Bemerkungen über Vorkommen und Lebensweise dieses Tieres. Engelmann, Leipzig 52 pp
- Schütz L (1961) Distribution and dispersal of the borer *Teredo navalis* L. and its entering of the Kiel Canal near Kiel. Kieler Meeresforschung 17: 228-236
- Schütz L (1963a) Die Beziehung zwischen Ei-, Embryonenzahl und Körpergröße der Weibchen einiger Peracaridae aus dem Nord-Ostsee-Kanal (Mesohalinikum). Zool. Anzeiger 171(5/8): 291-303
- Schütz L (1963b) Ökologische Untersuchungen über die Benthosfauna im Nordostseekanal. I: Autökologie der sessilen Arten. Int. Rev. ges. Hydrobiol. 48: 361-418
- Sikorski AV and Bick A (2004) Revision of *Marenzelleria* Mesnil, 1896 (Spionidae, Polychaeta). Sarsia 89: 253-275
- Sordyl H, Bönsch R, Gercken J, Gosselck F, Kreuzberg M and Schulze H (1998) Verbreitung und Reproduktion des Schiffsbohrwurms *Teredo navalis* L. an der Küste Mecklenburg-Vorpommerns. DGM 42(4): 142-149
- Souty-Grosset C, Holdich DM, Noel P, Reynolds JC and Hafner P (eds) (2006) Atlas of crayfish in Europe. Museum national d'Histoire naturelle, Paris (Patrimoines naturels, 64). p 187
- Spratte S and Hartmann U (1997) Fischartenkataster: Süßwasserfische und Neunaugen in Schleswig-Holstein. Ministerium Landwirtschaft, Ernährung und Tourismus, Kiel Germany, 183 pp
- Stachowicz JJ, Terwin JR, Whitlatch RB and Osman RW (2002) Linking climate change and biological invasions: Ocean warming facilitates nonindigenous species invasions. Proceedings - National Academy of Sciences, 99: 15497-15500
- Sukopp H and Brande A (1984) Beiträge zur Landschaftsgeschichte des Gebietes um den Tegeler See. Sitzungsber. Ges. Naturforsch. Freunde Berlin, 24: 198-214/1-7 pp
- Sures B and Streit B (2001) Eel parasite diversity and intermediate host abundance in the River Rhine, Germany. Parasitology 123: 185-191
- Szaniawska A and Dobrzycka-Krahel A (2004) National Report for Poland. ICES Working Group on Introductions and Transfers of Marine Organisms (WGITMO), Cesenatico Meeting 2004, 64-74
- Tambs-Lyche H (1964) *Gonionemus vertens* L. Agazziz (Limnomedusae) - a zoogeographical puzzle. Sarsia 15: 1-8
- Taraschewski H, Moravec F, Lamah T and Anders K (1987) Distribution and morphology of two helminths recently introduced into European eel populations: *Anguillicola crassus* (Nematoda, Dracunculoidea) and *Paratenuisentis ambiguus* (Acanthocephala, Tenuisentidae). Diseases of Aquatic Organisms 3: 167-176
- Thienemann A (1950) Verbreitungsgeschichte der Süßwassertierwelt Europas. Die Binnengewässer 18: 1-809
- Tittizer T (1996) Vorkommen und Ausbreitung aquatischer Neozoen (Makrozoobenthos) in den Bundeswasserstraßen. In: Gebhardt H, Kinzelbach R and Schmidt-Fischer S (eds.) Gebietsfremde Tierarten. Auswirkungen auf einheimische Arten, Lebensgemeinschaften und Biotope. Ecomed, Landsberg, 49-86

- Tittizer T, Schöll F, Banning M, Haybach A and Schleuter M (2000) Aquatische Neozoen im Makrozoobenthos der Binnenwasserstraßen Deutschlands. *Lauterbornia* 39: 1-72
- Tobias W (1972) Ist der Schlammröhrenwurm *Branchiura sowerbyi* BEDDARD 1882 (Oligochaeta: Tubificidae) ein tropischer Einwanderer im Untermain? *Natur und Museum* 102(3): 93-107
- Utting SD and Spencer BE (1992) Introductions of marine bivalve molluscs into the United Kingdom for commercial culture - Case histories. *ICES Mar. Sci. Symp.* 194, 84-91
- Van der Velde G, Rajagopal S, Kelleher B, Musko IB and bij de Vaate A (2000) Ecological impact of crustacean invaders: general considerations and examples from the Rhine River. In: von Vaupel Klein JC and Schramm FR (eds.) *The biodiversity crisis and Crustacea*. Proc. 4th Int. Crustacean Congress, Amsterdam, The Netherlands, July 20-24, 1998. Balkema, Rotterdam, 3-33
- Van Urk RM (1987) *Ensis americanus* (Binney) (syn. *E. directus* auct. non Conrad) a recent introduction from Atlantic North America. *J. Conchology* 32: 329-333
- Von Cosel R, Dörjes J and Mühlenhardt-Siegel U (1982) Die amerikanische Schwertmuschel *Ensis directus* (Conrad) in der Deutschen Bucht 1. Zoogeographie und Taxonomie im Vergleich mit dem einheimischen Schwertmuschel-Arten. *Senckenbergiana marit.* 14: 147-173
- Vranovsky M and Sporka F (1998) *Urnatella gracilis* Leidy 1851 (Kamptozoa) auch in der March. *Lauterbornia* 33: 85-93
- Vrieling EG, Koeman RPT, Nagasaki K, Ishida Y, Peperzak L, Gieskes WWC and Veenhuis M (1995) *Chattonella* and *Fibrocapsa* (Raphidophyceae): first observation of, potentially harmful, red tide organisms in Dutch coastal waters. *Neth. J. Sea Res.* 33: 183-191
- Wallentinus I (submitted) Chapter 2 Introductions and Transfers of Plants. Status of Introductions of Non-indigenous Marine Species to North Atlantic and adjacent Waters According to National Reports Considered at Meetings of the Working Group on introductions and Transfers of Marine Organisms 1992-2002. ICES Cooperational Research Report
- Wallentinus I (1992) Dispersal of *Sargassum muticum* in the ICES countries. Report of ICES WGITMO, 6 pp
- Wallentinus I (1999) Introduction and transfer of plants. In: Munro ALS, Utting SD and Wallentinus I (eds.) Status of introductions of non-indigenous marine species to North Atlantic waters 1981-1990. ICES Cooperative Research Report 231, 1-43
- Wawrzyniak-Wydrowska B and Gruszka P (2005) Population dynamics of alien gammarid species in the River Odra estuary. *Hydrobiologia*, 539: 13-25
- Weidema IR (2000) Introduced species in the Nordic countries. Nordic Council of Ministers, Copenhagen. Nord Environment, 2000:13, 242 pp.
- Welcomme RL (1988) International introductions of inland aquatic species. *FAO Fish. Tech. Pap.* 294: 318
- Werner B (1950) Die Meduse *Gonionemus murbachi* MAYER im Sylter Wattenmeer. *Zool. Jb.* 78: 471-505
- WGITMO (2005) Report of the Working Group on Introductions and Transfers of Marine Organisms. ICES CM 2005/ACME:05 Ref. E.: 173 pp. available at <http://www.ices.dk>
- WGITMO (2006) Report of the ICES Working Group on Introductions and Transfers of Marine Organisms. Oostende, Belgium. 234 pp. available at <http://www.ices.dk>
- Wiltshire KH and Dürselen C-D (2004) Revision and quality analysis of the Helgoland Reede long-term phytoplankton data archive. *Helgol. Mar. Res.* 58: 252-268
- Winkler HM, Skora K, Repecka R, Ploks M, Neelov A, Urho L, Gushin A and Jespersen H (2000) Checklist and status of fish species in the Baltic Sea. ICES CM 2000/Mini:11, 15 pp
- Wolff T (1977) The Horseshoe Crab (*Limulus polyphemus*) in North European waters. *Vidensk. Meddr Dansk Naturh. Foren.* 140: 39-52
- Wolff WJ (2005) Non-indigenous marine and estuarine species in The Netherlands. *Zoologische Mededelingen.* 79-1: 1-116
- Wunsch HH (1912) Eine neue Spezies des Genus *Corophium* Latr. aus dem Müggelsee bei Berlin. *Zool. Anz.* 39: 729-738
- Zibrowius H (1991) Ongoing modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Bull. Mus. Hist. Nat. Marseille* 51: 83-107

Checklist for aquatic alien species in Germany

**Annex**

Aquatic alien species reported from coastal areas of the North Sea and Baltic Sea and from inland waters within the national borders of Germany. Species which arrived by drift or other means of natural range expansion were not considered. IAS = an invasive alien species which threatens ecosystems, habitats or native species (sensu CBD 2000)

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<b>PHYTOPLANKTON</b>								
<b>Dinophyceae</b>								
<i>Karenia</i> (= <i>Gymnodinium mikimotoi</i> ) (Miyake et Kominami ex Oda) Hansen et Moestrup, 2000		1966		Pacific	unintentional	ships	bloom forming	Hickel et al. 1971
<i>Prorocentrum redfieldii</i> Bursa, 1959		<1999		unclear	unintentional	ships?	bloom forming	Nehring 1998b, Elbrächter 1999
<b>Raphidophyceae</b>								
<i>Chattonella antiqua</i> (Hada) Ono, 1980		1991		Pacific?	unintentional	Ships	potentially toxic	Elbrächter 1994, Vrieling et al. 1995, Lu and Göbel 2000
<i>Chattonella marina</i> (Subrahmanyam) Hara et Chihara, 1982		1991		Pacific?	unintentional	Ships	potentially toxic	Elbrächter 1994
<i>Fibrocapsa japonica</i> Toriumi et Takano, 1973		1991		Pacific?	unintentional	Ships	toxic / IAS	Elbrächter 1994
<b>Bacillariophyceae</b>								
<i>Coscinodiscus wailesii</i> Gran et Angst, 1931	1977	1977		Indo-Pacific	unintentional	aquaculture	competition, bloom forming / IAS	Hasle 1990, Laing 1999, Wiltshire & Dürselen 2004
<i>Odontella sinensis</i> (Greville) Grunow, 1884	1904	1903		Indo-Pacific	unintentional	Ships	competition	Ostenfeld 1908
<i>Thalassiosira punctigera</i> (Castracane) Hasle, 1983	<1983	1978		Indo-Pacific	unintentional	aquaculture	unknown	Hasle 1983, 1990
<b>MACROPHYTES</b>								
<b>Pteridophyta</b>								
<b>Azollaceae</b>								
<i>Azolla filiculoides</i> Lamarck, 1783			1980s	S America	intentional	ornamental trade	unknown	Hussner 2005
<b>Spermatophyta</b>								
<b>Apiaceae</b>								
<i>Hydrocotyle ranunculoides</i> Linnaeus, 1781			2004	N America	intentional	ornamental trade	competition, habitat modification / IAS	Hussner 2005
<b>Crassulaceae</b>								
<i>Crassula helmsii</i> (Kirk) Cockayne, 1907			1980s	Australia	intentional	ornamental trade	competition, habitat modification / IAS	Kowarik 2003, Hussner 2005
<b>Hydrocharitaceae</b>								
<i>Egeria densa</i> Planchon, 1849			1980s	S America	intentional	ornamental trade	unknown	Kowarik 2003, Hussner 2005
<i>Elodea canadensis</i> Michaux, 1803			1859	N America	intentional	ornamental trade	competition, habitat modification / IAS	Arndt 1931, Kowarik 2003, Hussner 2005, Wallentinus subm.
<i>Elodea nuttallii</i> (Planchon) St. John, 1920			1953	N America	intentional	ornamental trade	competition, habitat modification / IAS	Kowarik 2003, Hussner 2005, Wallentinus submitted

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<i>Vallisneria spiralis</i> Linnaeus, 1753 Lemnaceae			1966	N America	intentional	ornamental trade	unknown	Kowarik 2003, Hussner 2005
<i>Lemna minuta</i> Kunth, 1816			1983	N America	intentional	ornamental trade	unknown	Kowarik 2003
<i>Lemna turionifera</i> Landolt, 1975 Haloragaceae			1965	N America	intentional	ornamental trade	unknown	Kowarik 2003
<i>Myriophyllum aquaticum</i> (Velloso) Verdcourt, 1973			1980s	S America	intentional	ornamental trade	unknown	Hussner 2005
<i>Myriophyllum heterophyllum</i> Michaux, 1803 Poaceae			1962	N America	intentional	ornamental trade	unknown	Kowarik 2003
<i>Spartina anglica</i> Hubbard, 1968		1927		W Atlantic	intentional	planting	competition, habitat modification, hybridisation? / IAS	Kolumbe 1931, Dijkema 1983, Wallentinus submitted
Macroalgae Phaeophyceae								
<i>Ascophyllum nodosum</i> (Linnaeus) Le Jolis, 1863		1990s		unclear	unknown	unknown	unknown	Bartsch and Kuhlenkamp 2000, Wallentinus submitted
<i>Colpomenia peregrina</i> (Sauvageau) Hamel, 1937		1905		Pacific	unintentional	aquaculture	unknown	Fletcher and Farrell 1999
<i>Fucus evanescens</i> Agardh, 1820	1989			N Pacific	unintentional	ships	competition, habitat modification, hybridization	Wallentinus 1999, Hopkins 2001
<i>Sargassum muticum</i> (Yendo) Fensholt, 1955		1988		N Pacific	unintentional	aquaculture	fouling, habitat modification / IAS	Wallentinus 1992, Bartsch and Kuhlenkamp 2000, Wallentinus submitted
Rhodophyceae								
<i>Bonnemaisonia hamifera</i> Hariot, 1891		<1959		N Pacific	unintentional	aquaculture	competition	Kylin 1930, Bartsch and Kuhlenkamp 2000
<i>Dasya baillouviana</i> (Gmelin) Montagne, 1841	2002	1960s		W Atlantic	unintentional	aquaculture	unknown	Wallentinus pers. comm., Schories and Selig 2006
<i>Gracilaria vermiculophylla</i> (Ohmi) Papenfuss, 1967	2005	2002		Pacific	unintentional	aquaculture	unknown	Nehls 2004, Schories and Selig 2006
<i>Polysiphonia harveyi</i> Bailey, 1848		1960s		N Pacific	unintentional	aquaculture	unknown	Wallentinus 1999, Maggs and Stegenga 1999
Chlorophyceae								
<i>Codium fragile</i> ssp. <i>tomentosoides</i> (van Goor) Silva, 1955		1930s		N Pacific	unintentional	aquaculture	competition, fouling, habitat modification	Bartsch and Kuhlenkamp 2000, Wallentinus submitted
ZOOPLANKTON								
Ctenophora								
<i>Mnemiopsis leidyi</i> Agassiz, 1865	2006			unclear	unknown	unknown	unknown	Javidpour and Sommer pers. comm.

Checklist for aquatic alien species in Germany

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
Crustacea								
<i>Acartia tonsa</i> Dana, 1848	<1981	1931		Pacific / W Atlantic	unintentional	ships	competition	Klie 1933, Arndt and Schnese 1986
<i>Ameira divagans</i> Nicholls, 1939	1970s			W Atlantic	unintentional	ships	unknown	Scheibel 1974
<i>Cercopagis pengoi</i> (Ostroumov, 1891) MACROZOOBENTHOS	2004			Ponto-Caspian	unintentional	canal	competition, predation	Gruzka pers. com. in WGITMO 2005
Porifera								
<i>Eunapius carteri</i> Bowerbank, 1863			1993	Africa, Asia	unintentional	ornamental trade	fouling	Gugel 1995, Nehring 2002
Hydrozoa								
<i>Bimeria francisciana</i> Torrey, 1902		<1952	<1952	Indo-Pacific?	unintentional	ships	competition, habitat modification, predation	Schütz 1963a,b, Cohen and Carlton 1995
<i>Bougainvillia macloviana</i> Lesson, 1830		1895		Antarctic waters	unintentional	ships	unknown	Hartlaub 1897, Broch 1924
<i>Cordylophora caspia</i> (Pallas, 1771)	1870	1858	1899	Ponto-Caspian	unintentional	canal	competition, fouling, predation, habitat modification	Kirchenpauer 1862, Hinkelmann 1899, Schulze 1981, Gruszka 1999
<i>Craspedacusta sowerbyi</i> Lancaster, 1880			1923	NW Pacific	unintentional	ornamental trade	unknown	Tittizer 1996, Tittizer et al. 2000
<i>Gonionemus vertens</i> Agassiz, 1862		1947		N America	unintentional	ships	unknown	Werner 1950, Tams-Lyche 1964
<i>Nemopsis bachei</i> Agassiz, 1849		1942		W Atlantic	unintentional	ships	fouling?	Hartlaub 1911, Kühl 1962
Anthozoa								
<i>Cereus pedunculatus</i> (Pennant, 1777)		1921		NE Atlantic	unintentional	ships	unknown	Müllegger 1921, Pax 1936
<i>Diadumene cincta</i> (Stephenson, 1925)		1928		Pacific	unintentional	aquaculture	competition?	Pax 1936, Kluijver 1991
<i>Haliplanella luciae</i> (= <i>lineata</i> ) (Verrill, 1898)		1920		Pacific	unintentional	ships	fouling?	Pax 1920, Gollasch and Riemann-Zürneck 1996
Bivalvia								
<i>Congeria leucophaeta</i> (Conrad, 1831)	<1996	<1994	1928	N America	unintentional	ships	competition, fouling	Boettger 1933b, Post and Landmann 1994, Jungbluth 1996
<i>Corbicula fluminalis</i> (O.F. Müller, 1774)			1984	E Asia	unintentional	ships	competition / IAS	Kinzelbach 1991, Meister 1997, Nehring 2002
<i>Corbicula fluminea</i> (O.F. Müller, 1756)			1987	Asia	unintentional	ships	competition / IAS	Kinzelbach 1991, Nehring 2002
<i>Crassostrea angulata</i> (Lamarck, 1819)		1911		unclear	unintentional	aquaculture	unknown	Meyer-Waarden 1964
<i>Crassostrea gigas</i> (Thunberg, 1793)		1991		NW Pacific	unintentional	aquaculture	competition, habitat modification, hybridization ?, parasite carrier / IAS	Utting and Spencer 1992, Reise 1998a,b
<i>Crassostrea virginica</i> (Gmelin, 1791)	<1887	1911		unclear	unintentional	aquaculture	unknown	Möbius 1887, Rady 1913

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<i>Dreissena polymorpha</i> (Pallas, 1771)	1828	1835	1824	Ponto-Caspian	unintentional	canal	competition, fouling, habitat modification, parasite carrier / IAS	Dahl 1891, Bentheim-Jutting 1922, Arndt 1931, Thienemann 1950
<i>Ensis americanus</i> (Binney, 1870)	1993	1979		NW Atlantic	unintentional	ships	competition, habitat modification / IAS	Von Cosel et al. 1982, Essink 1985, van Urk 1987, Gürs et al. 1993
<i>Mya arenaria</i> (Linnaeus, 1758)	<1200	<1200	<1931	NW Atlantic	unintentional	ships	competition?	Arndt 1931, Petersen et al. 1992, Reise 1998a&b, Nehring 2000a
<i>Petricola pholadiformis</i> Lamarck, 1818	1927	1896		NW Atlantic	unintentional	aquaculture	competition, habitat modification	Schlesch 1932, Kuckuck 1957, Knudsen 1989, Jensen & Knudsen 2005
<i>Teredo navalis</i> Linnaeus, 1758	<1993	<1808		Indo-Pacific?	unintentional	ships	habitat modification / IAS	Hahn 1956, Schütz 1961, Sordyl et al. 1998
<i>Unio mancus</i> Lamarck, 1819 Gastropoda			<1922	S Europe	unintentional	canal	unknown	Tittizer et al. 2000
<i>Crepidula fornicata</i> (Linnaeus, 1758)		1934		W Atlantic	unintentional	aquaculture	competition, habitat modification, parasite carrier	Havinga 1929, Ankel 1935, Kuckuck 1957, Minchin et al 1995
<i>Gyraulus parvus</i> (Say, 1817)			1981	N America	intentional	ornamental trade	unknown	Geiter et al. 2002
<i>Lithoglyphus naticoides</i> (Pfeiffer, 1828)			1883	E Europe	unintentional	canal	parasite carrier	Thienemann 1950, Jungbluth 1996, Nehring 2002
<i>Menetus dilatatus</i> (Gould, 1841)			1980	N America	intentional	ornamental trade	unknown	Geiter et al. 2002
<i>Physella acuta</i> (Draparnaud, 1805)			1895	SW Europe	intentional	ornamental trade	unknown	Sukopp & Brande 1984, Jungbluth 1996, Nehring 2002
<i>Physella heterostropha</i> (Say, 1817)			<1927	N America	intentional	ornamental trade	unknown	Jungbluth 1996, Nehring 2002
<i>Planorbella duryi</i> (Weatherby, 1879)			1980s	N America	intentional	ornamental trade	unknown	Geiter et al. 2002
<i>Potamopyrgus antipodarum</i> (Gray, 1843) Platyhelminthes	1908		1900	New Zealand	unintentional	ships	competition, parasite carrier	Thienemann 1950, Cole 1982
<i>Dendrocoelum romano-danubiale</i> (Codreanu, 1949)			1992	Ponto-Caspian	unintentional	canal	unknown	Tittizer et al. 2000, Nehring 2002
<i>Dugesia tigrina</i> (Girard, 1850)			1931	N America	unintentional	ornamental trade	unknown	Hauer 1950, Tittizer 1996, Nehring 2002
Kamptozoa <i>Urnatella gracilis</i> Leidy, 1851			1960	N America	unintentional	ships	unknown	Franz 1992, Vranovsky and Sporka 1998, Geiter et al. 2002



Checklist for aquatic alien species in Germany

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<b>Oligochaeta</b>								
<i>Branchiura sowerbyi</i> Beddard, 1892			1959	W Pacific	unintentional	ornamental trade	habitat modification	Tobias 1972, Tittizer 1996, Gruszka 1999, Nehring 2002
<b>Polychaeta</b>								
<i>Aphelocheata marioni</i> (Saint-Joseph, 1894)		1938		unclear	unknown	unknown	unknown	Caspers 1950
<i>Ficopomatus enigmaticus</i> (Fauvel, 1923)			1975	S Pacific	unintentional	ships	fouling	Kühl 1977a, Cole 1982, Zibrowius 1991
<i>Hypania invalida</i> (Grube, 1860)			1995	Ponto-Caspian	unintentional	canal	habitat modification	Kothe 1968, Tittizer 1996, Nehring 2002
<i>Marenzelleria neglecta</i> (= cf. <i>viridis</i> ) Sikorski et Bick, 2004	1985	1996		NW Atlantic	unintentional	ships	competition, habitat modification, predation / IAS	Bick and Burckhardt 1989, Bick and Zettler 1997, Bastrop et al. 1997, Sikorski and Bick 2004
<i>Marenzelleria viridis</i> (= cf. <i>wireni</i> ) (Verrill, 1873)	2004	1983		NW Atlantic	unintentional	ships	competition, habitat modification, predation / IAS	Essink and Kleef 1986, Bick and Burckhardt 1989, Bick and Zettler 1997, Sikorski and Bick 2004
<i>Microphthalmus similis</i> Bobretzky, 1870		1962		unclear	unknown	unknown	unknown	Hartmann-Schröder and Stripp 1968
<i>Nereis virens</i> Sars, 1835	1920s	1923		unclear	unintentional?	ships?	predation	Reibisch 1926, Hagmeier and Kändler 1927, Hartmann-Schröder 1996
<i>Polydora ligérica</i> (Ferronière, 1898)			<1932	unclear	unknown	unknown	unknown	Augener 1940, Jaeckel 1962, Hartmann-Schröder 1996
<i>Tharyx killariensis</i> (Southern, 1914)		1972		unclear	unintentional?	aquaculture?	unknown	Hauser 1973, Hartmann-Schröder 1996
<b>Crustacea</b>								
<i>Astacus leptodactylus</i> Eschscholtz, 1823			1910s	Ponto-Caspian	intentional	stocking	unknown	Geiter et al. 2002, Souty-Grosset et al. 2006
<i>Atyaephyra desmarestii</i> (Millet, 1831)			1932	Mediterranean	unintentional	canal	unknown	De Lattin 1967, Tittizer 1996, Tittizer et al. 2000, Nehring 2002
<i>Balanus improvisus</i> Darwin, 1854	1867	1858	<1899	W Atlantic	unintentional	ships	fouling, habitat modification	Kirchenpauer 1862, Hoek 1875, Dechow 1920, Broch 1924, Bätke 1995
<i>Callinectes sapidus</i> Rathbun, 1896		1964		NW Atlantic	unintentional	ships	predation	Gruner 1962, Kühl 1965
<i>Caprella mutica</i> Schurin, 1935		2004		Pacific	unintentional	ships	clogging of gear?	Schrey and Buschbaum 2006

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<i>Chelicorophium curvispinum</i> (Sars, 1895)	1932	1920s	1912	Ponto-Caspian	unintentional	canal	competition, habitat modification / IAS	Wundsch 1912, Schlienz 1922, Neuhaus 1933, Tittizer 1996, Bernauer et al. 1996, Gruszka 1999, Nehring 2002
<i>Chelicorophium robustum</i> (Sars, 1895)			2004	Ponto-Caspian	unintentional	canal	unknown	Eggers and Martens 2004
<i>Corophium sextonae</i> Crawford, 1937		1997		S Pacific	unintentional	ships	competition	Nehring and Leuchs 1999
<i>Crangonyx pseudogracilis</i> Bousfield, 1958			1992	N America	unintentional	stocking?	unknown	Bernauer et al. 1996
<i>Dikerogammarus haemobaphes</i> (Eichwald, 1841)			1993	Ponto-Caspian	unintentional	canal	competition	Tittizer 1996, Nehring 2002, Wawrzyniak-Wydrowska and Gruszka 2005
<i>Dikerogammarus villosus</i> (Sovinsky, 1894)			1995	Ponto-Caspian	unintentional	canal	competition, predation / IAS	Tittizer 1996, Nehring 2002
<i>Echinogammarus berilloni</i> (Catta, 1878)			1924	Mediterranean	unintentional	canal	unknown	Tittizer 1996, Nehring 2002
<i>Echinogammarus ischnus</i> (Stebbing, 1899)			1977	Ponto-Caspian	unintentional	canal	unknown	Tittizer 1996, Tittizer 1996, Nehring 2002
<i>Echinogammarus trichiatus</i> (Martynov, 1932)			2000	Ponto-Caspian	unintentional	canal	unknown	Podraza et al. 2001, Nehring 2002
<i>Elminius modestus</i> Darwin, 1854		1953		S Pacific	unintentional	ships	competition?, fouling	Bishop 1947, Kühl 1954
<i>Eriocheir sinensis</i> Milne-Edwards, 1854	1932	1915	1912	NW Pacific	unintentional	ships	competition, habitat modification, parasite carrier, predation / IAS	Schnakenbeck 1924, Marquard 1926, Boettger 1933, Peters 1933, Peters and Hoppe 1938, Panning 1938
<i>Gammarus tigrinus</i> Sexton, 1939	1975	1965	1957	NW Atlantic	intentional	stocking	competition, parasite carrier, predation	Bousfield 1958, Schmitz 1960, Klein 1969, Bulnheim 1976, Nehring 2002, Wawrzyniak-Wydrowska and Gruszka 2005
<i>Hemimysis anomala</i> Sars, 1907			1997	Ponto-Caspian	unintentional	canal	competition, predation	Faasse 1998, Ketelaars et al. 1999, Nehring 2002
<i>Jaera istri</i> Vieuille, 1979			1995	Ponto-Caspian	unintentional	canal	unknown	Kothe 1968, Tittizer 1996, Nehring 2002
<i>Limnomysis benedeni</i> Czerniavsky, 1882			1997	Ponto-Caspian	unintentional	canal	unknown	Tittizer et al 2000, Nehring 2002
<i>Obesogammarus crassus</i> (Sars, 1894)			2004	Ponto-Caspian	unintentional	canal	unknown	Eggers and Anlauf 2005
<i>Obesogammarus obesus</i> (Sars, 1894)			2004	Ponto-Caspian	unintentional	canal	unknown	Nehring 2006c
<i>Orchestia cavimana</i> Heller, 1865			1920	Ponto-Caspian	unintentional	ships	unknown	Schlienz 1922, Tittizer 1996

Checklist for aquatic alien species in Germany

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<i>Orconectes immunis</i> (Hagen, 1870)			1997	N America	intentional	ornamental trade	parasite carrier, habitat modification / IAS	Geiter 1998, Souty-Grosset et al. 2006
<i>Orconectes limosus</i> (Rafinesque, 1817)			1890	NE America	intentional	stocking	parasite carrier, competition, predation / IAS	Schellenberg 1928, Boettger 1934, Sukopp and Brande 1984
<i>Pacifastacus leniusculus</i> (Dana, 1852)			1980s	N America	intentional	stocking	parasite carrier, competition, predation, habitat modification / IAS	Huber and Schubart 2005, Souty-Grosset et al. 2006
<i>Palaemon macrodactylus</i> Rathbun, 1902		2004		SE Asia	unintentional	ships	unknown	González-Ortegón et al. 2006
<i>Pontogammarus robustoides</i> (Sars, 1894)	1994		1994	Ponto-Caspian	unintentional	canal	competition, hybridization, predation	Rudolph 1997, Nehring 2002, Wawrzyniak-Wydrowska and Gruszka 2005
<i>Proasellus coxalis</i> (Dollfus, 1892)		<1987	1930s	Medi-terranean	unintentional	canal	unknown	Gruner 1965, Post and Landmann 1994, Tittizer 1996, Nehring 2002
<i>Proasellus meridianus</i> (Racovitza, 1919)			1930s	W Europa	unintentional	canal	unknown	Thienemann 1950, Gruner 1965, Tittizer 1996, Nehring 2002
<i>Procambarus clarkii</i> (Girard, 1852)			1990s	N America	intentional	ornamental trade	parasite carrier, competition, predation, habitat modification / IAS	Geiter et al. 2002, Souty-Grosset et al. 2006
<i>Rhithropanopeus harrisi</i> (Gould, 1841)	1936	<1977	1936	NW Atlantic	unintentional	ships	competition, predation	Neubaur 1936, Buitendijk and Holthuis 1949, Christiansen 1969, Van der Velde et al. 2000, Kühl 1977b, Cole 1982, Adema 1991, Tittizer 1996, Nehring 2000b
Chelicerata								
<i>Caspihalacarus hyrcanus</i> Vietz, 1928			<2006	Ponto-Caspian	unintentional	canal	unknown	Martens et al. 2006
<i>Limulus polyphemus</i> Linnaeus, 1758		1866		NW Atlantic	intentional	ornamental trade	unknown	Lloyd 1874, Holthuis 1950, Wolff 1977
Bryozoa								
<i>Pectinatella magnifica</i> (Leidy, 1851)			1883	N America	unintentional	ships	unknown	Tittizer et al. 2000, Nehring 2002
<i>Victorella pavid</i> Saville Kent, 1870	1911		1951	Indo-Pacific?	unintentional	ships	competition, habitat change	Kraeplin 1887, Ax 1952
Ascidiacea								
<i>Styela clava</i> Herdmann, 1882		1997		N Pacific	unintentional	ships	competition, fouling	Millar 1960, Reise 1998a,b

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
FISHES								
<i>Acipenser baerii</i> Brandt, 1869	1980s	1980s		Russia	intentional	stocking	hybridisation	Spratte and Hartmann 1997, Gessner et al. 1999
<i>Acipenser gueldenstaedti</i> Brandt et Ratzeberg, 1833	<1990s			unclear	intentional	stocking	hybridisation	Gerstmeier and Romig 1998
<i>Acipenser ruthenus</i> Linnaeus, 1758		<1992		unclear	intentional	stocking	hybridisation	Gerstmeier and Romig 1998, Gessner et al. 1999
<i>Acipenser transmontanus</i> Richardson, 1836		<1990s		N America	intentional	stocking	hybridisation	WGITMO 2006
<i>Ameiurus</i> (= <i>Ictalurus</i> ) <i>melas</i> (Rafinesque, 1820)			1990s	N America	intentional	stocking	competition, predation	Welcomme 1988, Spratte and Hartmann 1997
<i>Ameiurus</i> (= <i>Ictalurus</i> ) <i>nebulosus</i> (Lesueur, 1819)			1885	N America	intentional	stocking	competition	Spratte and Hartmann 1997
<i>Carassius auratus</i> (Linnaeus, 1758)			<1560	Asia	intentional	stocking	hybridization	Arnold 1990
<i>Coregonus peled</i> (Gmelin, 1789)			1965	Asia	intentional	stocking	hybridisation, predation	Geiter et al. 2002
<i>Lepomis cyanellus</i> Rafinesque, 1819			1965	N America	intentional	stocking	hybridisation	Arnold 1990
<i>Lepomis gibbosus</i> (Linnaeus, 1758)			1880	N America	intentional	stocking	competition, predation, hybridisation	Welcomme 1988, Spratte and Hartmann 1997
<i>Neogobius kessleri</i> (Günther, 1861)	<2004			unclear	unknon	unknown	competition, predation	Harka & Biro 2004
<i>Neogobius melanostomus</i> (Pallas, 1811)	1999			Ponto-Caspian	intentional	canal	competition, habitat modification, predation	Winkler at el. 2000, Szaniawska and Dobrzycka-Kraheil 2004
<i>Oncorhynchus mykiss</i> (Walbaum, 1792)			1882	N America	intentional	stocking	competition, habitat modification, hybridization, predation, parasite carrier	Welcomme 1988, Spratte and Hartmann 1997, Winkler at el. 2000
<i>Proterorhinus marmoratus</i> (Pallas, 1811)			1999	Ponto-Caspian	intentional	canal	unknown	Schadt 2000, Harka and Biro 2004
<i>Pseudorasbora parva</i> (Temminck et Schlegel, 1846)			1984	E Asia	intentional	stocking	competition	Spratte and Hartmann 1997
<i>Salvelinus fontinalis</i> (Mitchill, 1814)			1890	NW Atlantic	intentional	stocking	competition, hybridization, predation	Muus and Dahlström 1968
<i>Umbra krameri</i> Walbaum, 1792			<1997	unclear	intentional	ornamental trade	unknown	Spratte and Hartmann 1997
<i>Umbra pygmaea</i> (De Kay, 1842)		1924	1910s	N America	intentional	ornamental trade	competition	Duncker 1939, Spratte and Hartmann 1997
AMPHIBIAN								
Anura								
<i>Rana catesbeiana</i> Shaw, 1802			1990s	N America	intentional	ornamental trade	predation / IAS	Laufer 2004
PARASITES								
Oomycota								
<i>Aphanomyces astaci</i> Schikora, 1906			1878	N America	unintentional	stocking	crayfish parasite / IAS	Dehus 1990

Checklist for aquatic alien species in Germany

Species	Year of first record and recipient region			Origin / donor area	Vector	Pathway	Impact or potential impact / invasiveness	References
	Baltic Sea	North Sea	Inland waters					
<b>Acanthocephala</b>								
<i>Paratenuisentis ambiguus</i> (van Cleave, 1921)	?	?	1987	N America	unintentional	stocking	eel parasite	Taraschewski et al. 1987
<b>Platyhelminthes</b>								
<i>Pseudodactylogyrus anguillae</i> (Yin et Sproston, 1948)	?	?	1980s	E Asia	unintentional	stocking	eel parasite	Buchmann et al. 1987, Sures and Streit 2001
<i>Pseudodactylogyrus bini</i> (Kikuchi, 1929)	?	?	1980s	E Asia	unintentional	stocking	eel parasite	Buchmann et al. 1987, Sures and Streit 2001
<b>Nematoda</b>								
<i>Anguillicola crassus</i> (Kuwahara, Niimi et Hagaki, 1974)	1980s	1980s	1980s	E Asia	unintentional	stocking	eel parasite / IAS	Taraschewski et al. 1987, Minchin and Rosenthal 2002
<b>Annelida</b>								
<i>Barbronia weberi</i> (Blanchard, 1897)			1994	S Asia	unintentional	ornamental trade	predation	Potel et al. 1998
<i>Caspiobdella fadejewi</i> (Epshtein, 1961)			1990s	Ponto-Caspian	unintentional	canal	fish leech	Geissen and Schöll 1998
<i>Piscicola haranti</i> Jarry, 1960			1990s	Ponto-Caspian	unintentional	canal	fish leech	Schimmer 1995, Tittizer et al. 2000
<i>Xironogiton victoriensis</i> Gelder et Hall, 1990			2003	N America	unintentional	stocking	crayfish parasite	Martens et al. in press