

Rapid coastal survey for targeted alien species associated with floating pontoons in Ireland

Dan Minchin

Marine Organism Investigations, 3 Marina Village, Ballina, Killaloe, Co Clare, Ireland

E-mail: minchin@indigo.ie

Received 12 February 2007; accepted in revised form 22 February 2007

Abstract

Twenty alien species were targeted in a rapid assessment of all Irish coasts during 2005 to 2006 at twenty-nine floating pontoon sites consisting of marinas, jetties or fish-farms. Twelve species were found and the survey provided twenty-five new locality records. Four species had not been recorded previously in Ireland. Three of these were tunicates *Corella eumyota*, *Botrylloides violaceus* and a cryptogenic *Didemnum* sp. The fourth species was not targeted, the amphipod *Monocorophium insidiosum*. The majority of the new locality records (67%) were from the Irish Sea coast. Some of the targeted species were removed from leisure craft hulls: *Balanus improvisus*, *C. eumyota*, *Corophium mutica*, *Didemnum* sp. and *Styela clava*.

Key words: invasive species, marinas, non-natives, rapid assessment, Ireland

Introduction

Rapid assessment surveys assist management in controlling alien species. It may be possible to eliminate them when found at an early stage, reduce their rate of spread or develop techniques to mitigate effects to commercial interests. With few exceptions alien species arriving in Ireland normally first appear in Britain (Minchin and Eno 2002) and are usually found in port regions where merchant and fishing vessels berth, at marinas for leisure craft and in shellfish culture areas (Minchin 2006a, Minchin et al. 2006a). Pontoons of marinas and fish-farms were chosen in this study because the immersed parts of the floating pontoons are not usually treated with antifouling coatings and so can develop extensive plant and invertebrate growths and also may be easily sampled at all tidal states. Fish farm pontoons were used to provide an assessment in coastal areas where marina pontoons do not exist. Worldwide survey techniques vary in their intensity, their duration and purpose.

The port studies undertaken in Australia involved a wide range of sampling techniques, including diving, and these surveys provided extensive accounts of both native and alien biota (Hewitt et al. 1999). There is considerable effort involved in such surveys. In North America, pontoons at marina sites were chosen for sampling by specialists with expertise in different taxonomic groups (Pedersen et al. 2005). The relative ease that species can be collected make this approach time efficient. Many of the species collected in such surveys have little recognisable impact although some invasive forms are readily identifiable. Hayes et al. (2002) developed an approach involving a target list of species not present in Australia that could be sought in future surveys based on four criteria that included known invasiveness elsewhere, their presence with known vectors and whether these vectors were active. In their desk assessment, these were arranged according to relative risk of invasion.

In this risk assessment, features from each of these surveys have been used but these are dependant on the target species being readily

identifiable in the field. Such a survey was carried out in Scotland during 2006 (Ashton et al. 2006) where ten marina sites were surveyed revealing the presence of four out of seven targeted species. In this study twenty-nine sites were visited to search for ten species already introduced to Ireland and to determine whether a further ten other species had arrived in Ireland. Such an approach is likely to be of practical value in monitoring invasive species in the implementation of the European Union's Water Framework Directive and other management studies.

Methods

Twenty-nine marina or fish farm floating pontoons, from all Irish coasts (Figure 1 and see Annex 2 for positions), were sampled over the period March 2005 to August 2006. The majority of sites were visited during July and August 2006. Approximately one hour was spent at each site visiting different separate pontoon sections exposed to different conditions. Most marina sites had ca 20 to 100 floating units and other sites had 1 to ca 30 floating units. At each site the most sheltered part of the marina to the most exposed, to wind or current movements, were examined. Biota along the submerged sides and ends of pontoons as well as shaded areas where there were fewer macroalgae were examined. Samples were removed from pontoons from close to the water surface to depths of <1m and on the supporting poles for the pontoons to depths of 3.5m. Samples were collected using a scraper and pocket net on a pole or directly removed by hand.

At least six separate pontoon sections were visited at each site where this was possible. Heavily fouled boat hulls of up to 15m overall length berthed alongside marina boardwalks were also examined. Observation was greatly facilitated under sun-bright calm conditions. Selected samples were collected and directly preserved in 90% ethanol.

Target species were selected based on their current known distribution on the coasts of Ireland, Britain or Northern Europe (Annex 1). Twenty species were selected that could be sampled at three different levels: 1) easily recognised and/or sampled by hand from above the water surface 2) obtained with simple sampling equipment 3) easily removed from scraped samples to be later examined microscopically. Identification of unfamiliar species in advance of the survey was aided by a study of



Figure 1. Distribution of sampling sites, see also Annex 2 for positions

images obtained from websites and from photographs or specimens received from other workers. Voucher samples of new records to Ireland were collected and have been deposited in the National Museum of Ireland, Dublin.

Results and Discussion

Twelve of the twenty targeted species were found in this study. Two other non-targeted species, one alien and one cryptogenic, were also found. Four of the species were not previously known in Ireland. There were twenty-five new locality records.

Species not previously recorded in Ireland

Three tunicate species were found for the first time in Ireland. The extensive pendulous growths (Figure 2) of the *Didemnum* sp. are reported elsewhere in more detail (Minchin & Sides 2006). Colonies of *Didemnum* sp. were found at two marina sites on pontoons and leisure craft on the Irish Sea coast. It is likely to be a high impacting species of special concern to managers on account of it being able to rapidly foul surfaces. It may have impacts on mussels in hanging culture as a similar form has done in New Zealand (Coutts and Sinner 2004).

The southern circumpolar tunicate *Corella eumyota* was found at five sites on the south-west, southern and eastern Irish coasts. At all sites the collected specimens were in clusters or formed an extensive layer attached by their tunic. At Marloge marina in Cork Harbour it was the dominant fouling organism on the hull of a small yacht (Figure 3 and 4). Photographs of fouling organisms from Fenit taken in March 2005 show large specimens of *C. eumyota*, without any orange colouration, and was attached to *Styela clava*. The species was first recorded in Europe by Lambert (2003) from France and it is likely that it will rapidly expand its range further within Ireland.

The western Pacific colonial tunicate *Botrylloides violaceus* was found at Malahide and Carlingford marinas. The colonies are made up of different colour morphs. The yellow form was the most conspicuous and could be easily seen from a distance of over 10m. Colonies had grown over other fouling species such as mussels, solitary tunicates and algae (Figure 5). None appeared to have any fouling on their upper surfaces. This species is also known to foul shellfish installations (Cohen et al 2001). This can lead to increased labour in farming cultured molluscs.

The amphipod *Monocorophium insidiosum* (Crawford 1937) was found in the sheltered Carrickfergus marina on Belfast Lough. This species has not been recorded in Ireland before and was collected believing it to be the related *M. sextonae*. It is unclear whether the species is introduced. *M. sextonae* was also recorded in this survey and had previously been recorded in Ireland by Costello (1993). In some regions it has become abundant, such as in Lough Hyne on the Irish south-west coast, where it competes with native amphipods (JMC Holmes pers. comm.).

Species already recorded in Ireland

Extensions to the range of eight of the ten targeted species already present in Ireland were found. A further non-targeted species, the anemone *Haliplanella lineata* (Verrill, 1869), was found in Fenit in Tralee Bay. Four specimens were collected. While this bay has a wild oyster fishery, no layings of imported oysters are known. An introduction from such a source is possible. Individuals were found imported with half-grown oysters to Dungarvan Bay in 1993 and Waterford Harbour, since 2001, arriving from France. It is possible that it was introduced as hull fouling. Fenit is a small shipping port and ships are known



Figure 2. Pendulous growth of *Didemnum* sp. attached to a complete mussel shell, Malahide marina, January 2006 (Photo by Dan Minchin)



Figure 3. *Corella eumyota* (with orange siphons) from the hull of a yacht, Marloge marina, Cork Harbour, June 2006 (Photo by Dan Minchin)

to arrive here from northern Europe. The only other known locality in Ireland for *H. lineata* is from the Ardfry rapids ca 100km to the north in Galway Bay (Ryland and Nelson-Smith 1975).

The Pacific brown algae *Colpomenia peregrina* and *Sargassum muticum* are now present on most Irish coasts. *C. peregrina* had not been previously recorded from Schull Harbour and *S. muticum* was collected from three new localities. Algal fragments of *S. muticum* were found floating close to marina pontoons. Adjacent area searches revealed stands of the plant in shallow water or specimens cast ashore. Both species are dispersed naturally but some may be spread inadvertently with hull fouling or oyster movements.

At the sheltered marina at Arklow the serpulid *Ficopomatus enigmaticus* was found at depth on the supporting poles for the floating platform below a low salinity layer due to discharges of the Avoca River. *F. enigmaticus* was also found at the Poolbeg marina in the estuarine area of the River Liffey draining into Dublin Bay. This species was not previously known from the Irish east coast.

The crustacean *Monocorophium sextonae* continues to expand its range occurring on the southern and the eastern coasts but remains unknown on the western coast. In comparison, the Pacific amphipod *Caprella mutica* now occurs on east, south and west coasts. It was abundant in Dun Laoghaire Harbour during the summer but no specimens were found over-winter despite extensive sampling (Minchin and Holmes in press).

The tunicate *Styela clava* was found for the first time in the Owenabue Estuary at the Royal Cork Yacht Club, Crosshaven, Cork Harbour, attached to the hulls of two cabin cruisers one of which had been idle over two years. Specimens were found on hulls of craft in Marloge, Cork Harbour and Dun Laoghaire Harbour. While the species was known from the innermost part of Cork Harbour (Minchin and Duggan 1988), their presence at Crosshaven extends the known range within the harbour. With recent expansions to several areas in Ireland (Minchin et al. 2006b) it is likely to extend its range further. All present and previous collections and observations have been of small numbers. However, there are conditions when this species can cause extensive economic damage to the hanging culture of some molluscs (Bourque et al. 2003).

Species not found in the survey

The Pacific tunicate *Perophora japonica* was first described in Britain from the Fleet on the south coast (Baldock and Bishop 2001) and since then has appeared in south-west Wales (Bishop et al.

2003). The species is easily identifiable and may yet be found in Ireland.



Figure 4. Part of colony of *Corella eumyota* removed from a fouled yacht, Marloge marina, Cork Harbour, June 2006 (Photo by Dan Minchin)



Figure 5. Different colour morphs of *Botrylloides violaceus* overgrowing the mussel *Mytilus edulis* taken from the Carlingford marina, June 2006 (Photo by Dan Minchin)

The slipper limpet *Crepidula fornicata* has been recorded as being established in Ireland at one time but no populations are now known to exist. The species is present on the south coast of Britain, south-west and North Wales. Small males have been found in low numbers associated with imports of half-grown Pacific oysters from France arriving to Waterford Harbour since 2000. These will have been imported in mesh bags, none were found following some months of cultivation and had probably become crushed within the oysters bags. Small green crabs were regularly associated with imported oyster consignments and this may be a route for the entry of the Pacific grapsid crabs *Hemigrapsus penicillatis* and *H. sanguineus*. These crabs are normally found on

shores and it is unclear whether they are likely to be found on floating pontoons. Although *Crassostrea gigas* supports a large oyster fishery in Ireland, and apart from a small number of settled individuals being found in the wild, it is not known to recruit in Irish coastal waters. Nevertheless, there are concerns about large settlements elsewhere in northern Europe in The Netherlands and in Germany. Some settlements are also recorded from the south coast of Britain with rare settlements known from North Wales.

Chinese mitten crabs *Eriocheir sinensis* were found for the first time in Ireland in Waterford Harbour during 2006 (Minchin 2006b and J Caffrey pers. comm.). The nearest region where they are established is in the Thames River. Chinese mitten crabs are often recorded in small numbers in many north European ports. However, if introduced and established in Waterford Estuary the species may have consequences for inland fisheries. Since the time of this survey some more specimens have been collected in the same estuary (J Caffrey pers. comm.)

The north-west Pacific kelp *Undaria pinnatifida* is likely to arrive in Ireland but none were found on this occasion. It would be expected to be found attached to floating pontoons.

Assessment of the method

The survey method, by familiarisation of the species from photographs, was effective in finding targeted species that had not been seen before by a researcher. The survey revealed potential pests: *Didemnum* sp., *B. violaceus*, *C. eumyota*, *S. clava*, *F. enigmaticus*, and *S. muticum*. The majority of new distribution records were from the Irish Sea coast and it is likely that further species may be found at Carlingford. The method is likely to be more useful as more marinas are constructed. This method will provide for a better rapid coastal coverage and at low cost. It would seem practical that a larger number of species could be targeted in a survey of this nature. There is every possibility that some targeted species will have been overlooked; but if present in moderate abundance and not forming isolated clusters they would be expected to be seen.

The survey method has shortcomings. Although mobile species are found within the fouling biota (including decapoda, amphipoda, annellida, nematoda, copepoda, flatworms, and some fishes) the majority of the species found were sessile or adherent. The targeted crab species may not have been found in this study as these are more usually

found on shores or over sediments or other habitat types. Highly mobile species, such as fishes, are less likely to be found in such surveys. The use of grabs, baited traps and light traps are likely to reveal more alien species if used.

Acknowledgements

Field assistance was provided by Jon Williams and Barbara Minchin. I thank the marina and fish-farm managers for allowing access to sample. Photographs to aid field identification were kindly supplied by John Bishop and Gretchen Lambert and both confirmed the identification of collected tunicates and provided other useful information. I am especially indebted to Mark (JMC) Holmes of the national Museum who confirmed the identity of the crustaceans and for helpful discussions with J. Caffrey. The work forms part of a 'toolkit' funded by EU 6th Framework Projects Assessing Large-scale environmental risks with tested Methods (ALARM) contract number GOCE-CT-2003-506675 and presence and distributions of alien species was funded by Delivering Alien Invasive Species In Europe (DAISIE) contract number SSPI-CT-2003-511202.

References

- Ashton G, Boos K, Shucksmith R and Cook E (2006) Rapid assessment of the distribution of marine non-native species in marinas in Scotland. *Aquatic Invasions* 1 (4): 209-213
- Baldock B and Bishop JDD (2001) Occurrence of the non-native ascidian *Perophora japonica* in the Fleet, southern England. *Journal of the Marine Biological Association of the United Kingdom* 81: 1067
- Bishop JDD, Neal KJ, Nielsen S, Okawara B, Pemberton AJ and Wood C (2003) Biology of sessile invertebrates. *Marine Biological Association of the United Kingdom. Report to Council* 2003. 48 pp <http://www.mba.ac.uk/PDF/AR2003.pdf>
- Boaden PJS (1995) The adventive seaweed *Sargassum muticum* (Yendo) Fensholt in Strangford Lough, Northern Ireland. *Irish Naturalists' Journal* 25: 111-113
- Bourque D, Landry T, Davidson J, McNair N (2003) Impact of an invasive tunicate in Atlantic Canada: Recruitment and competition. *Journal of Shellfish Research* 22 (1): 320
- Clark PF, Rainbow PS, Robbins RS, Smith B, Yeomans WE, Thomas M and Dobson G (1998) The Alien Chinese mitten crab, *Eriocheir sinensis* (H. Milne Edwards, 1854) [Crustacea: Decapoda: Brachyura] in the Thames Catchment. *Journal of the Marine Biological Association*. 78(4): 1215-1221
- Cohen A, Berry AD, Milles CE, Milne D, Britton-Simmons K, Wonham MJ, Secord DL, Barks JA, Bingham B, Bookheim BE, Byers JE, Chapman JW, Cordell J, Dumbauld B, Fukuyama A, Harris LH, Kohn AJ, Li K, Mumfield jnr TF, Radashevsky V, Sewell AT and Welsch K (2001) Washington State exotics expedition 2000?: a

- rapid survey of exotic species in the shallow waters of Elliot Bay, Totten and Eld Inlets and Willapa Bay. Report to Washington State Department of Natural Resources, Olympia, Washington, USA. 52 pp
- Costello MJ (1993) Biogeography of alien amphipods occurring in Ireland and interactions with native species. *Crustaceana* 65: 287-299
- Coutts ADM and Sinner J (2004) An updated benefit-cost analysis of management options for *Didemnum vexillum* in Queen Charlotte Sound. Report to Marlborough District Council, New Zealand. Cawthron Report 925, 29pp http://woodshole.er.usgs.gov/projectpages/stellwagen/didemnum/images/pdf/news/cawthron_924.pdf
- Grave S, Moore SJ and Burnell G (1998) Changes in benthic macrofauna associated with intertidal oyster, *Crassostrea gigas* (Thunberg) culture. *Journal of Shellfish Research* 17: 1137-1142
- Eno NC, Clark RA and Sanderson WG (1997) Non-native species in British waters: a review and dictionary. Joint Nature Conservation Committee, Peterborough. 152 pp.
- Fletcher RL and Farrell P (1999) Introduced brown algae in the North East Atlantic, with particular respect to *Undaria pinnatifida* (Harvey) Suringar. *Helgoländer Meeresuntersuchungen* 52 (3-4): 259-275
- Gollasch S (1999) The Asian decapod *Hemigrapsus penicillatus* (de Haan, 1833) (Decapoda, Grapsidae) introduced in European waters, status quo and future perspective. *Helgoländer Meeresunters.* 52: 359-366
- Hayes KR, McEnulty FR and Silva C (2002) Identifying potential marine pests: an inductive approach. Centre for Research on Marine Pests. Report to Environment Australia. 42pp http://www.issg.org/database/species/reference_files/Hayes,%20K.R.%20et%20al.%202002.pdf
- Hewitt C, Campbell ML, Thresher RE and Martin RB (1999) Marine biological invasions of Port Phillip Bay, Victoria. Centre for Research on Introduced Marine Pests CSIRO Technical Report No. 20. 344 pp
- Lambert G (2003) New records of ascidians from the NE Pacific: a new species of *Trididemnum*, range extension and redescription of *Aplidiopsis pannosum* (Ritter, 1899) including its larva, and several non-indigenous species. *Zoosystema* 25: 665-679
- Minchin A (1991) Further distributional records of the adventive marine brown alga *Colpomenia peregrina* (Phaeophyta) in Ireland. *Irish Naturalists' Journal* 23: 38-381
- Minchin D (2004) Aquatic transport and the spread of aquatic species: challenges for management. In: Davenport J and Davenport JL (eds) *The effects of human transport on ecosystems: cars and planes, boats and trains*, 244-265. Dublin: Royal Irish Academy
- Minchin D (2006a) The transport and spread of living aquatic species. In: Davenport J and Davenport JD (eds) *The Ecology of Transportation: Managing Mobility for the Environment*. pp 77-97. Springer, The Netherlands.
- Minchin D (2006b). First record of the Chinese mitten crab *Eriocheir sinensis* (Milne-Edwards, 1854) (Decapoda: Crustacea). *Irish Naturalists' Journal* 28(7): 303-304
- Minchin D, Davis MH and Davis ME (2006b) Spread of the invasive tunicate *Styela clava* Herdman 1882 to the east and south-west coasts of Ireland. *Aquatic Invasions* 1(2): 91-96
- Minchin D, Duggan CB (1988) The distribution of the exotic ascidian, *Styela clava* Herdman, in Cork Harbour. *Irish Naturalists Journal* 22: 388-393
- Minchin D and Eno E (2002) Exotics of coastal and inland waters of Ireland and Britain. In: *Invasive Aquatic Species of Europe: Distribution, Impact and Management*. (eds) Leppäkoski E, Gollasch S and Olenin S. Kluwer, Dordrecht pp 267-275
- Minchin D, Floerl O, Savini D and Occhipinti-Ambrogi A (2006a) Small craft and the spread of exotic species. In: Davenport J and Davenport JD (eds) *The Ecology of Transportation: Managing Mobility for the Environment*. pp 99-118. Springer, The Netherlands
- Minchin D and Holmes JMC (in press). The first record of *Caprella mutica* Schurin 1935 (Crustacea: Amphipoda) from the east coast of Ireland. *Irish Naturalists' Journal*
- Minchin D, McGrath D and Duggan CB (1995) The slipper limpet *Crepidula fornicata* (L.) in Irish waters with a review of its occurrence in the north-eastern Atlantic. *Journal of Conchology* 35: 247-254
- Minchin D and Sides E (2006) Appearance of a cryptogenic tunicate, a *Didemnum* sp., fouling marina pontoons and leisure craft in Ireland. *Aquatic Invasions* 1(3): 143-147
- O'Riordan RM (1996) The current status and distribution of the Australian barnacle *Elminius modestus* Darwin in Ireland. In: Keegan BF and O'Connor R(eds) *Proceedings of the Irish Marine Science Symposium 1995*. Galway University Press, Galway pp 207-218
- Pedersen J, Bullock R, Carlton JT, Dijkstra J, Dobroski N, Dyrinda P, Fisher R, Harris L, Hobbs N, Lambert G, Lazo-Wasem E, Mathieson A, Miglietta M-P, Smith J, Smith III J and Tyrrell M (2005) Marine Invaders in the Northeast: Rapid assessment survey of non-native and native marine species of floating dock communities, August 2003. MIT Sea Grant College Program No 05-3, 46 pp
- Ryland JS and Nelson-Smith A (1975) Littoral and benthic investigations on the west coast of Ireland - IV (Section A: faunistic and ecological studies) Some shores in counties Clare and Galway. *Proceedings of the Royal Irish Academy* 75B: 245-266
- Tierney TD, Kane F, Naughton O, Kennedy S, O'Donohoe P, Copley L and Jackson D (2004) On the occurrence of the caprellid amphipod *Caprella mutica* Schurin 1935, in Ireland. *Irish Naturalists' Journal* 27: 437-439
- d'Udekem d'Acoz C and Faasse M (2002) De huidige status van *Hemigrapsus sanguineus* (de Haan, 1835) en *H. penicillatus* (de Haan, 1835) in de noordelijke Atlantische Oceaan, in het bijzonder in Nederland, met opmerkingen over hun biologie (Crustacea, Decapoda, Brachyura) [The current status of *Hemigrapsus sanguineus* (de Haan, 1835) and *H. penicillatus* (de Haan, 1835) in the northern Atlantic Ocean, with special emphasis on the Netherlands, with remarks on their biology]. *Het Zeepaard* 62(4): 101-115

Annex 1

Target list of species searched for during the 2005-2006 Irish survey.

Species	Sampling level	Nearest known region where established	Reference
<i>Colpomenia peregrina</i> Sauvageau, 1927	1	Ireland	Minchin 1991
<i>Undaria pinnatifida</i> (Harvey) Suringar, 1873	1	S coast of Britain	Fletcher & Farrell 1999
<i>Sargassum muticum</i> (Yendo) Fensholt, 1955	1	Ireland	Boaden 1995
<i>Balanus amphitrite</i> Darwin, 1854	2-3	S North Sea	Eno et al. 1997
<i>Balanus improvisus</i> Darwin, 1854	2-3	Ireland	Minchin 2004
<i>Botrylloides violaceus</i> Oka, 1927	1-2	S coast of Britain	JDD Bishop pers. comm.
<i>Caprella mutica</i> Schurin, 1935	1-3	Ireland	Tierney et al. 2004
<i>Corella eumyota</i> Traustedt, 1882	1-2	S coast of Britain	JDD Bishop pers. comm.
<i>Crassostrea gigas</i> Thunberg, 1793	2	Ireland	de Grave et al. 1998
<i>Crepidula fornicata</i> Linnaeus, 1758	2	Wales	Minchin et al. 1995
<i>Didemnum</i> sp	1	Northern France	Lambert pers. comm.
<i>Elminius modestus</i> Darwin, 1854	1	Ireland	O'Riordan 1996
<i>Eriocheir sinensis</i> H. Milne-Edwards, 1853	2	Thames River, Britain	Clark et al. 1998
<i>Hemigrapsus penicillatus</i> (de Haan, 1835)	2	Northern France and The Netherlands	Gollasch 1999
<i>Hemigrapsus sanguineus</i> (de Haan, 1853)	2	The Netherlands	d'Udekem d'Acoz and Faasse 2002
<i>Ficopomatus enigmaticus</i> (Fauvel, 1923)	1-3	Ireland	Minchin 2004
<i>Monocorophium sextonae</i> (Crawford, 1937)	2-3	Ireland	Costello 1993
<i>Perophora japonica</i> Oka, 1927	2	SW Wales	Bishop et al. 2003
<i>Rhithropanopeus harrisi</i> (Gould, 1841)	2	S Wales	D Minchin pers. ob. 2005
<i>Styela clava</i> Herdman, 1882	1-2	Ireland	Minchin et al. 2006b

Annex 2

Distribution of sampling sites in Ireland in 2005-2006 and records of non-native species found. New species records appear in bold

Map ref.	Location	Location coordinates		Dates of surveys	Species present	Dominant fouling organisms
		Latitude, °N	Longitude, °W			
1	Carlingford marina	54°03.01'	06°11.27'	03.08.2005 28.06.2006	<i>B. violaceus</i>, <i>C. mutica</i>, <i>C. eumyota</i>, <i>Didemnum</i> sp., <i>E. modestus</i>, <i>M. sextonae</i>, <i>S. muticum</i>	mussels, tunicates
2	Malahide marina	53°27.26'	06°09.21'	12.10.2006 05.07.2006	<i>B. violaceus</i>, <i>Didemnum</i> sp., <i>E. modestus</i>, <i>M. sextonae</i>	mussels
3	Howth marina	53°23.35'	06°03.68'	17.04.2005 18.01.2006 05.07.2006	<i>E. modestus</i>	tunicates
4	Poolbeg marina	53°20.40'	06°12.50'	05.07.2006	<i>F. enigmaticus</i>	barnacles tunicates
5	Dun Laoghaire marina	53°17.82'	06°08.08'	20.08.2005 18.01.2006 05.07.2006	<i>C. mutica</i>, <i>C. eumyota</i>, <i>M. sextonae</i>, <i>S. clava</i>, <i>E. modestus</i>	
6	Arklow marina	52°47.72'	06°08.66'	07.07.2006	<i>F. enigmaticus</i>, <i>E. modestus</i>	barnacles
7	Kilmore Quay marina	52°10.46'	06° 35.40'	07.07.2006	No alien species recorded	tunicates
8	Waterford marina	52°15.64'	07°06.28'	20.04.2006	<i>B. improvisus</i>	barnacles
9	Marloge marina,	51°52.02'	08°12.34'	17.06.2006	<i>C. mutica</i>, <i>C. eumyota</i>, <i>E. modestus</i>, <i>M. sextonae</i>, <i>S. clava</i>	mussels, tunicates
10	Royal Cork marina	51°48.28'	08°18.25'	21.06.2006	<i>E. modestus</i> , <i>M. sextonae</i>, <i>S. clava</i>	mussels
11	Crosshaven boatyard marina	51°48.32'	08°17.39'	21.06.2006	<i>E. modestus</i>	mussels
12	Kinsale marina	51°42.10'	08°31.05'	21.06.2006	<i>E. modestus</i>	mussels
13	Castlepark marina	51°41.78'	08°30.65'	21.06.2006	<i>E. modestus</i>	mussels
14	Schull pontoon	51°31.70'	09°32.89'	21.06.2006	<i>C. peregrina</i>, <i>E. modestus</i>	mussels
15	Cahirciveen marina	51°56.54'	10°14.04'	21.06.2006	No alien species recorded	mussels
16	Dingle Marina	52°08.29'	10°16.67'	20.06.2006	<i>E. modestus</i>	mussels tunicates
17	Fenit marina	52°16.25'	09°51.75'	31.03.2005 20.06.2006	<i>C. eumyota</i>, <i>E. modestus</i>, <i>S. clava</i>, <i>H. lineata</i>	
18	Kilrush marina	52°38.10'	09°29.67'	02.06.2005	<i>B. improvisus</i> , <i>F. enigmaticus</i>	varied
19	Rosroe pontoon	53°37.40'	09° 52.25'	03.07.2006	<i>E. modestus</i>	mussels
20	Mullaghmore pontoon	54°27.78'	08°26.23'	08.08.2005	No alien species recorded	mussels
21	Mulroy pontoon	55°13.38'	07°42.38'	29.06.2006	No alien species recorded	mussels
22	Swilly marina	55°05.13'	07°28.33'	29.06.2006	<i>E. modestus</i>	tunicates
23	Coleraine marina	55°07.60'	06°40.39'	29.06.2006	No alien species recorded	light fouling
22	Ballycastle marina	55°12.40'	06°16.38'	29.06.2006	No alien species recorded	mussels, tunicates
25	Carrickfergus marina	54°42.65'	05°48.80'	29.06.2006	<i>C. eumyota</i>, <i>M. insidiosum</i>	varied
26	Bangor marina	54°39.86'	05°40.19'	28.06.2006	<i>E. modestus</i>	mussels, tunicates
27	Donaghadee marina	54°39.10'	05°30.00'	28.06.2006	No alien species recorded	
28	Portaferry marina	54°22.57'	05°31.79'	28.06.2006	<i>S. muticum</i> , <i>E. modestus</i>	mussels
29	Ardglass marina	54°15.51'	05°36.50'	28.06.2006	<i>S. muticum</i>, <i>E. modestus</i>	mussels