

## ***Dikerogammarus villosus* (Sowinski, 1894) in the River Odra estuary – another invader threatening Baltic Sea coastal lagoons**

Piotr Gruszka<sup>1\*</sup> and Adam Woźniczka<sup>2</sup>

<sup>1</sup>Agricultural University in Szczecin, Department of Marine Ecology and Environmental Protection, Królewicza Kazimierza 4, 71-550 Szczecin, Poland

<sup>2</sup>Sea Fisheries Institute in Gdynia, Research Station in Świnoujście, Plac Słowiański 11, 72-600 Świnoujście, Poland

\*Corresponding author

E-mail: [pgruszka@fish.ar.szczecin.pl](mailto:pgruszka@fish.ar.szczecin.pl)

Received 29 July 2008; accepted in revised form 14 October 2008; published online 18 December 2008

### **Abstract**

*Dikerogammarus villosus* is a pontogammarid species that has rapidly extended its range in Europe starting from the Ponto-Caspian basin. In the 1990s it moved from the Hungarian to Austrian section of the River Danube, and then (after the opening of the Main-Danube canal) invaded German inland waterways. Using the canals joining the different river systems, the species has reached the River Odra. There it quickly spread both up- and downstream, reaching for the first time the Baltic Sea coastal waters (the Szczecin Lagoon) by 2002, extending the list of alien species that can be found in the Baltic. Based on samples taken in 2002-2004, *D. villosus* occurrence in the Szczecin Lagoon and adjacent coastal waters were identified. A mass occurrence of the species was found in some places already in 2002. Owing to the fact that an important (both for open sea and inland traffic) shipping route crosses the lagoon, populated by the new species, it is very probable that the area can act as a stepping stone in dispersal of *D. villosus* to other oligohaline Baltic Sea coastal waters.

**Key words:** *Dikerogammarus villosus*, Szczecin Lagoon, introduced species; coastal lagoons; colonization

### **Introduction**

Baltic coastal waters harbour at present all of the already established non-indigenous species that have made their way into the Baltic; in the coastal zone, those species are abundant and dominant (Nikolayev 1974; Leppäkoski and Olenin 2001; Leppäkoski et al. 2002). Baltic estuaries and coastal lagoons seem to offer particularly favourable conditions for those species (Olenin and Leppäkoski 1999; Orlova et al.

1999; Ezhova et al. 2005; Schories and Selig 2006). Therefore, the biological diversity in those areas is greatly affected by the presence of non-indigenous species; their distribution has been extended due to human mediation and has been termed xenodiversity (Leppäkoski and Olenin 2000). The River Odra (Oder) estuary, with the Szczecin Lagoon in its central part, is one of the xenodiversity centres in the Baltic (Gruszka 1999; Leppäkoski et al. 2002). As the major harbours in the area are meeting points for seaways and inland waterways, the estuary has

been invaded by both brackish and freshwater immigrant species which became permanent, and in some cases dominant, components of the biota, particularly in the macrozoobenthos (Masłowski 1992; Osadczuk and Wawrzyniak-Wydrowska 1998; Gruszka 1999; Rödiger 2004). Wide tolerance to changes in, e.g., salinity, has made it possible for some benthic species such as *Dreissena polymorpha* (Pallas, 1771), *Potamo-pyrgus antipodarum* (Grey, 1843), *Chelicorophium curvispinum* (G. O. Sars, 1895), and *Cordylophora caspia* (Pallas, 1771) to broaden their range and to settle in estuarine areas of central and western Europe, including Baltic coastal waters (Nikolayev 1951; Jazdzewski 1980; Eno et al. 1997; Leppäkoski and Olenin 2000). By the beginning of the 20th century, those species managed to colonise the Szczecin Lagoon (see Gruszka 1999).

Recently, however, more newly immigrated malacostracans, mostly of Ponto-Caspian origin, have spread spectacularly over Europe, often invading estuarine areas, including those in the Baltic Sea (Olenin and Leppäkoski 1999; Nehring 2000; Jazdzewski and Konopacka 2002; Zettler 2002; Arbačiauskas 2005; Grabowski et al. 2005; Jazdzewski et al. 2005; Berezina 2007; Grabowski et al. 2007a). The spread has been facilitated by connections of river basins via man-made canals and by intentional introductions (Jazdzewski 1980; Tittizer 1996; Bij de Vaate et al. 2002). Among those newcomers, gammarid amphipods are the dominant group so far with the following 8 species reported from the Baltic coastal waters: *Chaetogammarus ischnus* (Stebbing, 1899), *C. warpachowskyi* (G.O. Sars, 1894), *Dikerogammarus haemobaphes* (Eichwald, 1841), *D. villosus* (Sowinski, 1894), *Gammarus tigrinus* Sexton, 1939, *Gmelinoides fasciatus* (Stebbing, 1899), *Obesogammarus crassus* (G.O. Sars, 1894), and *Pontogammarus robustoides* (G.O. Sars, 1894).

So far *D. villosus*, a species belonging to the Pontogammaridae (a family originating from the Ponto-Caspian region), has been recorded only from the Szczecin Lagoon. The species has rapidly extended its range in Europe: migrating upstream the Danube, it spread from its native Black Sea basin (Carauşu et al. 1955; Mordukhai-Boltovskoi 1979; Neesemann et al. 1995) to the Slovak-Hungarian and then, in the 1980s and early 1990s, to the Austrian and German sections of the Danube (Csanyi 1994; Neesemann et al. 1995). The species may co-occur with two other congeners: *D. bispinosus*


(Martynov, 1925 (Müller et al. 2002; Kley and Maier 2006) and *D. haemobaphes* (Jazdzewski and Konopacka 1988; Kley and Maier 2006; this paper). *D. bispinosus* was earlier regarded as a subspecies of *D. villosus* (e.g. Carauşu et al. 1955; Mordukhai-Boltovskoi et al. 1969), but now, after Barnard and Barnard (1983), is referred to as a separate species (e.g. Jazdzewski and Konopacka 1988), the status being supported by the findings of genetic studies carried out by Müller and Schramm (2001) and Müller et al. (2002). According to these authors, the species reached the upper Danube (the Austrian waters) by 1998, and is expected to extend its range into the North Sea basin, as was earlier the case with (a) *D. haemobaphes* in 1993, in two successive waves of invasion (Schleuter et al. 1994; Schöll et al. 1995; Leuchs and Schleuter 1996) and (b) *D. villosus* in 1994 (Bij de Vaate and Klink 1995; Schöll et al. 1995). The expansion of those two species was possible due to the opening of the Main-Danube canal in 1992. Having invaded the River Rhein system, *D. villosus* could have displaced *D. haemobaphes* in Germany (Van der Velde et al. 2000, Kley and Maier 2006) and moved farther to colonise Swiss, French, and Dutch inland waterways (Devin et al. 2001; Bij de Vaate et al. 2002; Josens et al. 2005; Wittenberg 2006). The species moved eastwards by using canals joining different river systems in northern Germany (Grabow et al. 1998; Zettler 1999; Rudolph 2002), and reached the River Odra/Oder in the Baltic Sea basin between 1998 and 1999 (Müller et al. 2001; Jazdzewski and Konopacka 2002; Müller and Hertel 2004). Once in the Odra, it quickly spread both up- and downstream to reach the river's estuarine system. Thus, by appearing in the brackish Szczecin Lagoon by 2002, it made its first entry into the Baltic Sea coastal waters.

As described by Gruszka et al. (2003), samples collected in 2001 and 2002 showed the species to occur, in 2002, in the downstream reaches of the Odra south of Szczecin and in the southern part of the lagoon itself. Referring to their unpublished data, Jazdzewski et al. (2005), too, mentioned finding few individuals of *D. villosus* in samples collected from the lagoon in 2004, and thus confirmed the presence of the new invader in the Odra estuary.

In this paper, we discuss the distribution of *D. villosus* in the Szczecin Lagoon and its adjacent areas (Lake Dąbie and the terminal downstream section of the Odra) to summarise the first records of the species in Baltic coastal waters.

## Study area

As described by Majewski (1980) and other authors (e.g., Robakiewicz 1993; Osadczyk and Wawrzyniak-Wydrowska 1998), the Szczecin Lagoon (German: Stettiner Haff or Oderhaff; Polish: Zalew Szczeciński) is a shallow (3.8 m mean depth), 910 km<sup>2</sup> area that forms the core part of the River Odra (Oder) estuary (Figure 1).



**Figure 1.** The Szczecin Lagoon and adjacent waters (A.: Achterwasser)

The lagoon is influenced both by the riverine discharge from the Odra and smaller rivers (e.g., the Peene and the Uecker) and by inflows of Baltic waters from the Pomeranian Bay; as a result, its salinity is about 1-2 PSU (Majewski 1980; Landsberg-Uzciwek 2004). The bottom in the nearshore zone and in the Odra mouth is sandy, mud being the prevalent sediment type of the central part. Shores are covered by littoral vegetation, dominated mainly by phytocoenoses of *Phragmitetum australis* (Gams 1927) Schmale 1939 (Majewski 1980). The lagoon is geographically divided into a smaller part to the west, belonging to Germany (Kleines Haff), and

a larger, eastern part belonging to Poland (Wielki Zalew). This study was carried out in the Odra mouth (including the deltaic, freshwater Lake Dąbie) and in the Polish part of the lagoon.

## Materials and Methods

The sites supporting *D. villosus* in the Polish part of the Szczecin Lagoon and in the Odra downstream reaches up to the city of Szczecin were identified based on qualitative samples collected in 2002-2004. Figure 2 and Annex 1 show the sites sampled in particular years. The samples were obtained with various techniques used to collect benthic crustaceans, including a dredge (with 1 mm mesh), a Van Veen grab (0.7 mm sieve was used to extract the fauna), and a hand sieve (2 mm). In 2002, amphipods were also picked directly from stones retrieved from the shallow bottom and placed on a sheet of linen. Information on sampling techniques used at sites where *D. villosus* was found is given in Annex 1. The sampled animals were fixed in 10 % formalin. Amphipod collections from a total of 40 sites were checked for the presence of *D. villosus*.

## Results


Annex 1 summarises the data produced by analysing samples of amphipods collected from the Odra north of Szczecin and from shallow waters in the Polish part of the Szczecin Lagoon. Three samples collected in 2002 (Figure 2) revealed the presence of *D. villosus* in the studied section of the river (location 1) and in the southern area of the lagoon in the vicinity of the Odra mouth (locations 2 and 3). In 2003, the species was found in the northern part of the lagoon on a shallow sandy bottom covered with stones, within *Dreissena* clumps at the foot of the cliff shore south of the Island of Wolin off Lubin (location 4); and at two other sites (locations 5 and 6) situated along the southern shores of the lagoon (Figure 2). The 2004 survey confirmed the results obtained in the preceding years (Figure 2) to show the presence of the invader both south of the lagoon, i.e., in the eastern part of Lake Dąbie (locations 10 and 11), in the Odra off Police (location 7), and in the Odra mouth area north of Police (locations 8 and 9). The species was found also to be present in the southern part of the lagoon at the Polish-

German border (location 14) as well as in its northernmost part - in Lake Wicko and south of the Island of Wolin cliff in Lubin (locations 12 and 13).

The species was found to be accompanied by other alien amphipods that had earlier invaded the River Odra estuary (Annex 1). These are (in the order of the species' appearance): *Chelicorophium curvispinum*, *Gammarus tigrinus*, *Pontogammarus robustoides*, *Obesogammarus crassus*, and *D. haemobaphes*. Additionally, another alien amphipod, *Chaetogammarus ischnus*, rare in River Odra, was for the first time found in the Odra estuary, at a location invaded by *D. villosus* (Annex 1). No native gammarid amphipods were present in the samples containing *D. villosus*. *D. villosus* was not found in rivers Świna and Dziwna (sampled in 2002 and 2004), the watercourses connecting the Szczecin Lagoon with the Baltic Sea (the Pomeranian Bay).

## Discussion

Having invaded the River Odra estuary at the end of the 20th century, the alien gammarid amphipods *Gammarus tigrinus*, *Pontogammarus robustoides*, and *Obesogammarus crassus* became permanent and often dominant components of the Szczecin Lagoon macrozoobenthos (Gruszka 1999; Konopacka 2003; Rödiger 2004; Jądzewski et al. 2005; Wawrzyniak-Wydrowska and Gruszka 2005). The native *Gammarus pulex* (L., 1758), reported from the area until the 1970s (see Jądzewski and Konopacka 1995), retreated from the Lagoon, probably - as suggested earlier by Gruszka (1999) - due to an increase in salinity recorded in the area in the 1980s (Landsberg-Ucziwek 2001; Radziejewska and Chabior 2004). However, according to Konopacka (2003), a possibility that the elimination of the native gammarids was facilitated by the presence of the two newcomers (notably *G. tigrinus* and *P. robustoides*) cannot be ruled out. Schmid (1999) related the low abundance of *G. pulex* in the Lower Odra to the presence of *G. tigrinus*. The species native to the Baltic, mainly *G. duebeni* Liljeborg, 1852 and *G. zaddachi* Sexton, 1912, can be found in areas affected by inflows of the Baltic waters in the northern part of the Szczecin Lagoon and in the straits connecting the Lagoon with the Pomeranian Bay (Gruszka 1999; Gruszka, unpublished data; Jądzewski et al. 2005).



**Figure 2.** Location of sampling sites in the Szczecin Lagoon and adjacent waters sampled in 2002-2004. Empty frames show sites where *D. villosus* was not found, sites supporting *D. villosus* are marked with numbers corresponding to those given in Table 1: 1-3 (for 2002), 4-6 (for 2003), and 7-14 (for 2004). Details are provided in Annex 1

*D. villosus* and *D. haemobaphes* are two other newcomers that have appeared in the Odra estuary almost at the same time. The arrival of both species must have been quite recent, i.e., at the very beginning of the 21st century. As *D. villosus* was present in the samples collected in summer 2002, it can be inferred that this species reached the Lagoon as early as in 2001. About one year earlier, the two *Dikerogammarus* species, having successfully spread throughout Europe, were found in the downstream waters of the Odra: *D. haemobaphes* arrived from the east, while *D. villosus* came from the west (Müller et al. 2001; Bij de Vaate et al. 2002; Jądzewski and Konopacka 2002; Gruszka et al. 2003; Grabowski et al. 2007a). Once in the downstream Odra, *D. villosus* very soon gained

dominance over *D. haemobaphes* which, only a year earlier, was more abundant than its relative that had come from the west (Müller and Hertel 2004). In addition, *D. villosus* achieved dominance over other alien gammarid amphipods (*G. tigrinus* and *P. robustoides*) which had earlier spread upstream in the Odra (Müller et al. 2001; Müller and Hertel 2004). Further quantitative research should reveal whether these scenarios will be repeated in the Szczecin Lagoon.

The competitive potential of *D. villosus* was described in a number of studies: the larger *D. villosus* (the species can grow up to 3 cm; Neseemann et al. 1995) displaced *D. haemobaphes* at most locations in the upper Danube and in the Rhine system, thus stopping its further dispersal in southern Germany (Weinzierl et al. 1996; Haas et al. 2002; Kley and Maier 2006). It was shown to outcompete both native and alien gammarids in the Rhine and its deltaic system (Haas et al. 2002; Van der Velde et al. 2002; van Riel et al. 2007). Regarding the competition with *D. haemobaphes*, Kley and Maier (2006) suggest that *D. villosus* acts as a more effective predator. It is also known that, owing to its predatory behaviour, *D. villosus* is able to eliminate other gammarids, both native (*Gammarus duebeni*, *G. pulex*) and alien (*G. tigrinus*), by preying on them (Dick and Platvoet 2000; Haas et al. 2002; MacNeil and Platvoet 2005; van Riel et al. 2006). Large adult *D. villosus* are capable of attacking other invertebrates, e.g. *Asellus aquaticus*, *Chelicorophium curvispinum*, and some insects. As shown by stable isotope studies, the large predatory adults occupy the trophic level identical to that of some fish (Dick et al. 2002; van Riel et al. 2006). However, *D. villosus* is an omnivore (van Riel et al. 2006) capable of feeding also on microalgae (Platvoet et al. 2006). A broad diet spectrum or omnivory is only one of a set of attributes of successful aquatic invaders, as described by Ricciardi and Rasmussen (1998) and by Van der Velde et al. (2000). Other attributes include a short life span and a short generation time, a rapid growth with an early sexual maturity, a larger size than that of most relatives, a high reproductive capacity, a high genetic variability, a wide environmental tolerance, gregariousness, and an ability to disperse rapidly in association with human activities (e.g. shipping). *D. villosus* was shown to possess a high reproductive capacity, manifested in the very high fecundity; comparatively small eggs; small juveniles compared to adult body

length, which implies rapid growth; early maturity; a short generation time; and multiple reproduction (three generations per year) (Devin et al. 2004; Kley and Maier 2006; Grabowski et al. 2007b; Pöckl 2007). Comparison of these life history traits among different (both native and invasive) amphipod species made by Grabowski et al. (2007b) indicates *D. villosus* as belonging to the group of species with high competitive potential.

Devin et al. (2004) pointed to the female-biased sex-ratio as another characteristic that can favour successful establishment of the species in new regions. Neseemann et al. (1995) and Reinhold and Tittizer (1997) clearly demonstrated the role of ships in spreading the gammarid along inland waterways. Bruijs et al. (2001) and Wijnhoven et al. (2003) showed that *D. villosus* is an eurytopic species with a wide range of tolerance to temperature and salinity, a trait that renders the species capable of colonising the eutrophic Baltic Sea coastal lagoons. These lagoons are well prepared to receive a new invader that prefers hard substrates formed by clumps of the earlier Ponto-Caspian invader, *Dreissena polymorpha*, similarly to the situation in the Rhine (Haas et al. 2002). Another Baltic coastal water body, the Vistula Lagoon, will probably be soon colonised by *D. villosus* as the species has already entered the River Vistula basin: it was reported from River Bug in 2003 (Konopacka 2004). When this happens, the probability of the species' spreading into other coastal waters of the Baltic will increase significantly. This in turn will augment the risk of the species' invading the Great Lakes of North America (Ricciardi and Rasmussen 1998; Bruijs et al. 2001).

## Conclusions

*D. villosus*, a potential new invader, reached the Szczecin Lagoon by 2002 and become another constituent of the amphipod assemblages in some shallow bottom locations covered by stones and/or *D. polymorpha*. Monitoring of the adjacent waters should show whether *D. villosus* is in future capable of colonising the lagoon and adapting to varying salinity, typical of the Świna and the northernmost section of the Dziwna. Because the Szczecin Lagoon is intersected by a shipping route important both for ocean-going and for coastal and inland ship traffic, it is very probable that the area may play a role of a

stepping stone in the species' dispersal into other oligohaline coastal waters of the Baltic Sea.

### Acknowledgements

The authors are grateful to Dr Teresa Radziejewska for linguistic assistance. Our thanks are also due to two anonymous referees for their comments that helped to improve the final version of the manuscript.

### References

- Arbačiauskas K (2005) The distribution and local dispersal of Ponto-Caspian peracarida in Lithuanian fresh waters with notes on *Pontogammarus robustoides* population establishment, abundance and impact. *Oceanological and Hydrobiological Studies* 34 (Suppl. 1): 93-111
- Barnard JL, Barnard CM (1983) *Freshwater Amphipoda of the World*. Hayfield Association, Mt. Vernon, Virginia, 830 pp.
- Berezina N (2007) Invasions of alien amphipods (Amphipoda: Gammaridea) in aquatic ecosystems of North-Western Russia: pathways and consequences. *Hydrobiologia* 590: 15-29
- Bij de Vaate A, Klink AG (1995) *Dikerogammarus villosus* Sowinsky (Crustacea: Gammaridae) a new immigrant in the Dutch part of the Lower Rhine. *Lauterbornia* 20: 51-54
- Bij de Vaate A, Jazdzewski K, Ketelaars HAM, Gollasch S, Van der Velde G (2002) Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in Europe. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1159-1174
- Brujjs MCM, Kelleher B, van der Velde G, Bij de Vaate A (2001) Oxygen consumption, temperature and salinity tolerance of the invasive amphipod *Dikerogammarus villosus*: Indicators of further dispersal via ballast water transport. *Archiv für Hydrobiologie* 152: 633-646
- Caraușu S, Dobrea E, Manolache C (1955) Amphipoda of Brackish and Fresh Waters. *Fauna Republici populare Romine, Crustacea*, 409 pp.
- Csanyi B (1994) The macrozoobenthon community of the Danube between Rajka and Budapest. *Miscelanea Zoologica Hungarica* 9: 105-116
- Devin S, Beisel J-N, Bachmann V, Moreteau JC (2001) *Dikerogammarus villosus* Amphipoda: Gammaridae: Another invasive species newly established in the Moselle River and French hydrosystems. *Annls. Limnol.* - *Int. J. Limnol.* 37: 21-27
- Devin S, Piscart C, Beisel J-N, Moreteau JC (2004) Life history traits of the invader *Dikerogammarus villosus* (Crustacea: Amphipoda) in the Moselle River, France. *International Review of Hydrobiology* 89: 21-34
- Dick JTA, Platvoet D (2000) Invading predatory crustacean *Dikerogammarus villosus* eliminates both native and exotic species. *Proceedings of the Royal Society, Series B: Biological Sciences* 267: 977-983
- Dick JTA, Platvoet D, Kelly DW (2002) Predatory impact of the freshwater invader *Dikerogammarus villosus* (Crustacea: Amphipoda). *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1078-1084
- Eno NC, Clark RA, Sanderson WG (eds) (1997) *Non-native marine species in British waters: a review and directory*. Joint Nature Conservation Committee, Peterborough, 152 pp.
- Ezhova E, Żmudzinski L, Maciejewska K (2005) Long-term trends in the macrozoobentos of the Vistula Lagoon, southeastern Baltic Sea. Species composition and biomass distribution. *Bulletin of the Sea Fisheries Institute* 164(1): 55-73
- Grabow K, Eggers TO, Martens A (1998) *Dikerogammarus villosus* Sowinsky (Crustacea: Amphipoda) in norddeutschen Kanälen und Flüssen. *Lauterbornia* 33: 103-107
- Grabowski M, Jazdzewski K, Konopacka A (2005) Alien Crustacea in Polish waters – Introduction and Decapoda. *Oceanological and Hydrobiological Studies* 34 (Suppl. 1): 43-61
- Grabowski M, Jazdzewski K, Konopacka A (2007a) Alien Crustacea in Polish waters – Amphipoda. *Aquatic Invasions* 2(1): 25-38
- Grabowski M, Bacela K, Konopacka A (2007b) How to be an invasive gammarid (Amphipoda: Gammaroidea) – comparison of life history traits. *Hydrobiologia* 590: 75-84
- Gruszka P (1999) The River Odra Estuary as a Gateway for Alien Species Immigration to the Baltic Sea Basin. *Acta hydrochimica et hydrobiologica* 27(5): 374-382
- Gruszka P, Wawrzyniak-Wydrowska B, Żurawska J (2003) Alien crustacean species in the river Odra estuary (Baltic Sea). In: *Baltic Sea Science Congress 2003*, Helsinki, Finland, August 24-28, 2003. Abstract Publication, p. 130
- Haas G, Brunke M, Strei B (2002) Fast turnover in dominance of exotic species in the Rhine River determines biodiversity and ecosystem function: an affair between amphipods and mussels. In: *Leppäkoski E, Gollasch S, Olenin S (eds) Invasive Aquatic Species of Europe: Distribution, Impacts and Management*. Kluwer Academic Publishers, Dordrecht, pp 426-432
- Jazdzewski K (1980) Range extension of some gammaridean species in European inland waters caused by human activity. *Crustaceana Suppl.* 6: 84-107
- Jazdzewski K, Konopacka A (1988) Notes on the gammaridean amphipoda of the Dniester River Basin and Eastern Carpatians. *Crustaceana Suppl.* 13: 72-89
- Jazdzewski K, Konopacka A (1995) *Pancerzowce prócz równonogów lądowych Malacostraca prócz Oniscoidea*. (Malacostracans, except for Oniscoidea). *Katalog fauny Polski Catalogus faunae Poloniae [XIII]*, 1, Polska Akademia Nauk, Muzeum i Instytut Zoologii, Warszawa, 165 pp.
- Jazdzewski K, Konopacka A (2002) Invasive Ponto-Caspian species in waters of the Vistula and Oder basins and the southern Baltic Sea. In: *Leppäkoski E, Gollasch S and Olenin S (eds) Invasive Aquatic Species of Europe: Distribution, Impacts and Management*. Kluwer Academic Publishers, Dordrecht, pp. 384-398
- Jazdzewski K, Konopacka A, Grabowski M (2005) Native and alien malacostracan Crustacea along the Polish Baltic Sea coast in the 20th century. *Oceanological and Hydrobiological Studies* 34 (Suppl. 1): 175-193
- Josens G, Bij de Vaate A, Usseglio-Polatera P, Cammaerts R, Cherot F, Grisez F, Verboonen P, Vanden Bossche J-P (2005) Native and exotic Amphipoda and other Peracarida in the River Meuse: new assemblages emerge from a fast changing fauna. *Hydrobiologia* 542: 203-220

- Kley A, Maier G (2006) Reproductive characteristics of invasive gammarids in the Rhine-Main-Danube catchment, South Germany. *Limnologica* 36: 79-90
- Konopacka A (2003) Further step to the west - *Obesogammarus crassus* (G. O. Sars, 1894) (Crustacea, Amphipoda) already in the Szczecin Lagoon. *Lauterbornia* 48: 67-72
- Konopacka A (2004) Inwazyjne skorupiaki obunogie (Crustacea, Amphipoda) w wodach Polski. *Przegląd Zoologiczny XLVIII*, 3-4 (2004): 141-162
- Landsberg-Uczciwek M (ed) (2001) Raport o stanie środowiska w województwie zachodniopomorskim w roku 2000. IOŚ Biblioteka Monitoringu Środowiska, Szczecin, 199 pp.
- Landsberg-Uczciwek M (ed) (2004) Raport o stanie środowiska w województwie zachodniopomorskim w latach 2002-2003. IOŚ Biblioteka Monitoringu Środowiska, Szczecin, 336 pp.
- Leppäkoski E, Olenin S (2000) Non-native species and rates of spread: lessons from the brackish Baltic Sea. *Biol. Inv.* 2: 151-163
- Leppäkoski E, Olenin S (2001) The Meltdown of Biogeographical Peculiarities of the Baltic Sea: The Interaction of Natural and Man-made Processes. *Ambio* 30 (4-5): 202-209
- Leppäkoski E, Gollasch S, Gruszka P, Ojaveer H, Olenin S, Panov V (2002) The Baltic - a sea of invaders. *Canadian Journal of Fisheries and Aquatic Sciences* 59: 1175-1188
- Leuchs H, Schleuter A (1996) *Dikerogammarus haemobaphes* (Eichwald 1841), eine aus der Donau stammende Kleinkrebsart (Gammaridae) im Neckar. *Lauterbornia* 25: 139-141
- MacNeil C, Platvoet D (2005) The predatory impact of the freshwater invader *Dikerogammarus villosus* on native *Gammarus pulex* (Crustacea: Amphipoda); influences of differential microdistribution and food resources. *Journal of Zoology (London)* 267: 31-38
- Majewski A (ed) (1980) Zalew Szczeciński. WKiL, Warszawa, 339 pp.
- Masłowski J (1992) Bottom macrofauna of the Szczecin Lagoon (north-western Poland). *Acta Hydrobiologica* 34 (3): 253-274
- Mordukhai-Boltovskoi FD (1979) Composition and Distribution of Caspian Fauna in the Light of Modern Data. *International Revue ges. Hydrobiol.* 64 (1): 1-38
- Mordukhai-Boltovskoi FD, Greze II, Vasilenko SV (1969) Otrjad amfipody, ili raznogie - Amphipoda LATREILLE, 1816-1817. In: Vodjanickij VA (ed) *Opredelitel fauny Chernogo i Azovskogo morej*. Izdatelstvo "Naukova Dumka", Kiev, pp 440-524
- Müller JC, Schramm S (2001) A third *Dikerogammarus* invader is located in front of Vienna. *Lauterbornia* 41: 49-52
- Müller JC, Schramm S, Seitz A (2002) Genetic and morphological differentiation of *Dikerogammarus* invaders and their invasion history in Central Europe. *Freshwater Biology* 47: 2039-2048
- Müller O, Hertel A (2004) Abundanzentwicklung der invasiven Amphipoda *Dikerogammarus villosus* (Sovinski 1894) und *D. cf. haemobaphes* (Eichwald 1841) in der deutschen Oder und den angrenzenden Kanälen (Crustacea; Amphipoda). *Schriftenreihe des Bundesministeriums für Verbraucherschutz, Ernährung und Landwirtschaft (BMVEL)*, Reihe A: *Angewandte Wissenschaft* 498: 245-249
- Müller O, Zettler ML, Gruszka P (2001) Verbreitung und Status von *Dikerogammarus villosus* (Sovinski 1894) (Crustacea: Amphipoda) in der mittleren und unteren Strom-Oder und den angrenzenden Wasserstraßen. *Lauterbornia* 41: 105-112
- Nehring S (2000) Neozoen im Makrozoobenthos der deutschen Ostseeküste. *Lauterbornia* 39: 117-126
- Nesemann H, Pöckl M, Wittmann KJ (1995) Distribution of epigeal Malacostraca in the middle and upper Danube (Hungary, Austria, Germany). *Miscelanea Zoologica Hungarica* 10: 49-68
- Nikolayev II (1951) On new introductions from distant areas in fauna and flora of the North and the Baltic Seas. *Zoologicheskij Zhurnal* 30: 556-561
- Nikolayev II (1974) Principal tendencies in the biology of the recent Baltic Sea. *Oceanology (Moscow)* 16 (2): 1059-1069
- Olenin S, Leppäkoski E (1999) Non-native animals in the Baltic Sea: alteration of benthic habitats in coastal inlets and lagoons. *Hydrobiologia* 393: 233-243
- Orlova MI, Panov VE, Krylov PI, Telesh IV, Khlebovich VV (1999) Changes in planktonic and benthic communities of the eastern part of the Gulf of Finland (Baltic Sea) due to biological invasions. *Russian Academy of Sciences. Proc. Zool. Inst. St. Petersburg*, 279: 305-325
- Osadczyk A, Wawrzyniak-Wydrowska B (1998) Sediments in the Szczecin Lagoon: selected elements and macrozoobenthos. *Greifswalder Geographische Arbeiten* 16: 169-193
- Platvoet D, Dick JTA, Konijnendijk N, Van der Velde G (2006) Feeding on micro-algae in the invasive Ponto-Caspian amphipod *Dikerogammarus villosus* (Sowinsky, 1894). *Aquatic Ecology* 40(2): 237-245
- Pöckl M (2007) Strategies of a successful new invader in European fresh waters: fecundity and reproductive potential of the Ponto-Caspian amphipod *Dikerogammarus villosus* in the Austrian Danube, compared with the indigenous *Gammarus fossarum* and *G. roeseli*. *Freshwater Biology* 52: 50-63
- Radziejewska T, Chabior M (2004) Climatic and hydrological controls over the zoobenthos in a southern Baltic coastal lagoon. *Hydrobiologia* 514: 171-181
- Reinhold M and Tittizer T (1997) Zur Rolle von Schiffen als Vektoren beim Faunenaustausch Rhein/Main/Main-Donau-Kanal/Donau. *Deutsche gewässerkundliche Mitteilungen, Koblenz* 41: 199-205
- Ricciardi A, Rasmussen JB (1998) Predicting the identity and impact of future biological invaders: a priority for aquatic resource management. *Canadian Journal of Fisheries and Aquatic Sciences* 55: 1759-1765
- Robakiewicz W (ed) (1993) Warunki hydrodynamiczne Zalewu Szczecińskiego i cieśnin łączących Zalew z Zatoką Pomorską. (Hydrodynamics of the Szczecin Lagoon and straits connecting the Lagoon with the Pomeranian Bay). *Instytut Budownictwa Wodnego Polskiej Akademii Nauk, Gdańsk, Bibl. Nauk Hydrotechn.*, 16, 287 pp.
- Rödiger S (2004) Die Makrofauna des Oderhaffs - Vorschläge für ein Monitoring nach EU-WRRL (The macroinvertebrate community of the Oder Lagoon - Proposals for a monitoring after EU-WFD). In: Schernewski G and Dolch T (eds) *The Oder Estuary - against the background of the European Water Framework Directive*. *Marine Science Reports* 57, pp 127-178
- Rudolph K (2002) Über Veränderung der malakostraken Krebsfauna von Berlin und Brandenburg mit Angaben zum aktuellen Stand der Verbreitung. *Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin (N. F.)* 41: 93-108

- Schleuter M, Schleuter A, Potel S, Banning M (1994) *Dikerogammarus haemobaphes* (Eichwald 1841) (Gammaridae) aus der Donau erreicht ueber den Main-Donau-Kanal den Main. *Lauterbornia* 19: 155-159
- Schmid U (1999) Zur Besiedlung der Flußaue des Unteren Odertals durch Amphipoda (Crustacea). *Limnologie aktuell* Bd 9: 353-367
- Schöll F, Becker C, Tittizer T (1995) Das Makrozoobenthos des schiffbaren Rheins von Basel bis Emmerich 1986-1995. *Lauterbornia* 21: 115-137
- Schories D, Selig U (2006) Die Bedeutung eingeschleppter Arten (alien species) für die Europäische Wasserrahmenrichtlinie am Beispiel der Ostsee. *Rostock. Meeresbiol. Beitr.* 15: 147-158
- Tittizer T (1996) Vorkommen und Ausbreitung aquatischer Neozoen in den europäischen Bundeswasserstrassen. In: Gebhardt H, Kinzelbach R and Schmidt-Fischer S (eds) *Gebietsfremde Tierarten. Auswirkungen auf einheimischen Arten, Lebensgemeinschaften und Biotope. Situationsanalyse.* Umweltministerium Baden Württemberg. Ecomed Verlagsgesellschaft, Landsberg, pp 49-86
- Van Riel MC, Van der Velde G, Rajagopal S, Marguillier S, Dehairs F, Bij de Vaate A (2006) Trophic relationships in the Lower Rhine food web during invasion and after establishment of the Ponto-Caspian invader *Dikerogammarus villosus*. *Hydrobiologia* 565: 39-58
- Van Riel MC, Healy EP, Van der Velde G, Bij de Vaate A (2007) Interference competition among native and invader amphipods. *Acta Oecologica* 31: 282-289
- Van der Velde G, Rajagopal S, Kelleher B, Musko IB, Bij de Vaate A (2000) Ecological impact of crustacean invaders: General considerations and examples from the Rhine river. In: von Vaupel Klein JC and Schram FR (eds) *The Biodiversity crisis and Crustacea.* - Proceedings of the fourth international crustacean congress, Amsterdam, Netherlands, 20-24 July 1998, 2, pp. 3-33
- Van der Velde G, Nagelkerken I, Rajagopal S, Bij de Vaate A (2002) Invasions by alien species in inland freshwater bodies in Western Europe: The Rhine Delta. In: Leppäkoski E, Gollasch S and Olenin S (eds) *Invasive Aquatic Species of Europe: Distribution, Impacts and Management.* Kluwer Academic Publishers, Dordrecht, pp 360-372
- Wawrzyniak-Wydrowska B, Gruszka P (2005) Population dynamics of alien gammarid species in the River Odra estuary. *Hydrobiologia* 539: 13-25
- Weinzierl A, Potel S, Banning M (1996) *Obesogammarus obesus* (Sars 1894) in der oberen Donau (Amphipoda, Gammaridae). *Lauterbornia* 26: 87-89
- Wijnhoven S, van Riel MC, van der Velde G (2003) Exotic and indigenous freshwater gammarid species: physiological tolerance to water temperature in relation to ionic content of the water. *Aquatic Ecology* 37: 151-158
- Wittenberg R (2006) Crustaceans - Crustacea. In: Wittenberg R (ed.) *An inventory of alien species and their threat to biodiversity and economy in Switzerland.* CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forests and Landscape. The environment in practice no. 0629. Federal Office for the Environment, Bern, pp 65-70
- Zettler ML (1999) Erstnachweis von *Dikerogammarus villosus* (Sovinski, 1894) und Wiederfund von *Gammarus varsoviensis* Jażdżewski, 1975 in Mecklenburg-Vorpommern (Crustacea: Amphipoda). *Archiv der Freunde der Naturgeschichte in Mecklenburg* 38: 231-233
- Zettler ML (2002) Crustaceologische Neuigkeiten aus Mecklenburg-Vorpommern. *Archiv der Freunde der Naturgeschichte in Mecklenburg* 41: 15-35

**Annex 1**

Records of *Dikerogammarus villosus* in the Polish part of the Szczecin Lagoon and adjacent waters in 2002-2004 with sampling techniques indicated (D, dredg.; G, grab; H, hand siev; S, picking from stones onto a sheet of linen)

Year	No.	Geographic coordinates		Record date	Depth (m)	No. of <i>D. villosus</i> individuals collected	Other amphipod species found in the location	Habitat description (sampling technique)
		Latitude, N	Longitude, E					
2002	1.	53° 33.25'	14° 37.06'	4.07.2002	0.5	> 1000	<i>C. ischnus</i> , <i>D. haemobaphes</i> , <i>O. crassus</i> , <i>P. robustoides</i>	stones on sandy bottom, bulrushes and reeds (S, H)
	2.	53° 39.40'	14° 31.64'	11.07.2002	0.5	> 100	<i>G. tigrinus</i> , <i>D. haemobaphes</i> , <i>O. crassus</i> , <i>P. robustoides</i> , <i>C. curvispinum</i>	stones on sandy bottom, roots of trees (S)
	3.	53° 40.30'	14° 32.05'	3.10.2002	1	1	<i>P. robustoides</i> , <i>G. tigrinus</i> , <i>O. crassus</i>	sandy bottom (D)
2003	4.	53° 51.78'	14° 26.49'	2.07.2003	0.7	25	<i>G. tigrinus</i> , <i>O. crassus</i> , <i>P. robustoides</i> , <i>C. curvispinum</i>	sandy bottom with pebbles/stones, clumps of <i>Dreissena polymorpha</i> (G)
	5.	53° 42.25'	14° 21.45'	2.10.2003	0.5	10	<i>G. tigrinus</i> , <i>O. crassus</i> , <i>P. robustoides</i> , <i>C. curvispinum</i>	reeds (G)
	6.	53° 40.67'	14° 25.11'	2.10.2003	0.5	13	<i>G. tigrinus</i> , <i>O. crassus</i> , <i>P. robustoides</i>	reeds (G)
2004	7.	53° 33.40'	14° 35.75'	27.03.2004	2.5	3	<i>G. tigrinus</i> , <i>D. haemobaphes</i>	sandy bottom with pebbles/stones, clumps of <i>Dreissena polymorpha</i> (D)
	8.	53° 38.95'	14° 36.44'	2.08.2004	1.5	8	<i>G. tigrinus</i> , <i>D. haemobaphes</i>	bulrushes, sticking, organic sediment, covered with shells and sand (G)
	9.	53° 39.95'	14° 34.50'	6.08.2004	0.5	14	<i>D. haemobaphes</i> , <i>P. robustoides</i> , <i>C. curvispinum</i>	clump of <i>D. polymorpha</i> on the shell of <i>Anodonta</i> from sandy bottom (H)
	10.	53° 27.73'	14° 42.33'	2.09.2004	2	6	<i>D. haemobaphes</i>	clumps of <i>D. polymorpha</i> , sponges (G)
	11.	53° 27.73'	14° 42.41'	2.09.2004	1.5	16	<i>D. haemobaphes</i> , <i>O. crassus</i> , <i>P. robustoides</i>	gravel, densely packed sand, pebbles, <i>D. polymorpha</i> (G)
	12.	53° 53.37'	14° 25.35'	2.7.2004	0.5	9	<i>G. tigrinus</i>	sandy-gravelly bottom with stones (G)
	13.	53° 52.26'	14° 25.37'	25.10.2004	1.5	1	<i>G. tigrinus</i> , <i>O. crassus</i>	sand, bulrushes (G)
	14.	53° 44.39'	14° 17.59'	8.7.2004	0.7	3	<i>G. tigrinus</i>	dense reed belt (G)