

Research Article

Potential of the invasive colonial ascidian, *Didemnum vexillum*, to limit escape response of the sea scallop, *Placopecten magellanicus*

Jennifer A. Dijkstra^{1*} and Riley Nolan²¹Wells National Estuarine Research Reserve, 342 Laudholm Farm Road, Wells, Maine 04090 USA²University of New Hampshire, 46 College Road, Durham, NH 03824, USAE-mail: dijkstra@wellsnerr.org (JAD), rcu7@unh.edu (RN)

*Corresponding author

Received: 22 November 2010 / Accepted: 14 January 2011 / Published online: 14 July 2011

Editor's note:

This paper is a contribution to the proceedings of the 3rd International Invasive Sea Squirt Conference held in Woods Hole, Massachusetts, USA, on 26–28 April 2010. The conference provided a venue for the exchange of information on the biogeography, ecology, genetics, impacts, risk assessment and management of invasive tunicates worldwide.

Abstract

Didemnum vexillum is a globally invasive species and a major pest to the aquaculture industry. Like other colonial ascidians, *D. vexillum* can readily overgrow aquaculture nets and cultured species. Recently, the species has been found in great abundance on seafloor habitats, where it is often associated with commercially important shellfish species such as sea scallops, *Placopecten magellanicus*. Despite the increasing abundance of *D. vexillum* in areas that are regularly fished for sea scallops, little work has been conducted on the ascidians impact on scallop behavior. This study examined the effect of overgrowth of the sea scallop by *D. vexillum* using four measures: time to initial exhaustion, swimming speed, horizontal and vertical displacement. Scallops covered by *D. vexillum* became exhausted more quickly, and were not able to swim as far in either the horizontal or vertical direction as the control sea scallops without *D. vexillum* encrustation. The expansion of *D. vexillum* into sea scallop habitat may increase the vulnerability of sea scallops to predation and limit their ability to access food rich habitats.

Key words: escape response, epibiont, *Didemnum vexillum*, *Placopecten magellanicus*, swimming behavior

Introduction

The sea scallop, *Placopecten magellanicus* (Gmelin, 1791), is indigenous to the Northwest Atlantic, ranging from waters off Newfoundland and Labrador to Cape Hatteras, North Carolina at depths between 15 and 110 m (Posgay 1957). This fishery represents one of the largest fishing industries in the United States with a catch of 24,000 tonnes that is valued at over \$400 million US and is one of the most valuable wild scallop fisheries in the world (NMFS 2007). Sea scallop habitat and fishery extend both offshore and inshore along the eastcoast of the USA within the federal 3-mile Exclusive Economic Zone. In recent years, scallop landings in the 3-mile zone have declined, particularly along the coast of Maine (NMFS 2007), and likely would have declined on Georges Bank but for area closures

in the 1990s which resulted in a rebound in populations when fishing pressure was removed (Hart and Rago 2006). The preferred habitat of *P. magellanicus* is coarse sediment such as cobble, gravel, and clay (Hart and Chute 2004). In recent years, the sea scallop has been associated with and found to be encrusted by the colonial sea-squirt *Didemnum vexillum* Kott, 2002 (Valentine et al. 2007). While most non-native species occur in shallow waters on artificial and rocky substrate (Ruiz et al. 1997; Dijkstra et al. 2007; Neves et al. 2007), *D. vexillum* is often found in deeper waters in habitat preferred by sea scallops (Valentine et al. 2007; Mercer et al. 2009).

Scallops are bivalves and are often associated with encrusting epibionts including sponges, barnacles, polychaete worms, ascidians and bryozoans. However, the interactions between

scallops and different epibionts vary depending upon the identity of the encrusting species. For example, some sponges have a mutualistic relationship with scallops, by serving as protection against predation by sea stars either through camouflage or by interfering with the adhesion of the sea star's tubefeet (Forester 1979). Sponges also benefit by greater food intake (e.g., phytoplankton and/or bacteria) resulting from increased current generated by the scallop siphons (Forester 1979), or by being cleared of sediment when the scallop swims or claps its valves (Burns and Bingham 2002). In contrast, barnacles limit the ability of the scallop to swim long distances by decreasing the speed of the scallop and the swimming height it can attain, but confer no apparent benefit on the scallop (Donovan et al. 2003).

Swimming is an important escape response, particularly in sea scallops that are < 80 mm shell height (Baird 1954), allowing scallops to avoid predation and to disperse. Inhibiting the swimming behavior of scallops may increase the likelihood of predation (Stephens and Boyle 1978; Peterson et al. 1982). They also swim to migrate and to access habitats richer in food (Moore and Marshall 1967; Morton 1980; Hamilton and Koch 1996); thus, detrimental shell encrustation may limit their dispersal ability.

The expansion of *Didemnum vexillum* into valuable sea scallop fishing grounds in New England has led to concern about the impact of this tunicate on economically important sea scallop habitat (Valentine et al. 2007). One study by Morris et al. (2009) found that *D. vexillum* inhibits the settlement of the bay scallop (*Argopecten irradians irradians*), and therefore, will likely prevent the settlement of sea scallop spat. However, it is unclear whether *D. vexillum* affects the swimming behavior of sea scallops and if so to what degree. Therefore, we experimentally examined the effects of epibiosis by *D. vexillum* on the swimming behavior of *Placopecten magellanicus*. Given the importance of commercial shellfisheries, it is imperative to understand further ecological interactions between *D. vexillum* and juvenile and adult sea scallops.

Materials and methods

Our experiments were carried out in a flow-through sea-water clear Plexiglas tank (3.0 m × 0.85 m × 1.4 m) at the Shoals Marine Laboratory

on Appledore Island, ME in May and June, 2008. To quantify swimming behavior of *Placopecten magellanicus*, a plastic sheet spanning the length of the tank and marked with vertical lines 2 cm apart was attached to the wall of the tank. Another sheet of plastic (0.24 m × 1.4 m), also marked by 2 cm increments, was placed vertically in the tank. Swimming behavior of sea scallops was recorded using a video recorder mounted on a tripod and was placed ~1 meter in front of the tank. Videos were later analyzed for vertical and horizontal distance swum, initial time to exhaustion, and swimming speed.

To examine the effects of overgrowth by *Didemnum vexillum* on the escape response of *Placopecten magellanicus*, two treatments [no epibionts (control) and epibionts] and four response variables (time to exhaustion, swimming speed, horizontal distance, and vertical distance) were measured. A total of 38 scallops with shell heights ranging from 30 mm to 70 mm were used for the experiment. In each of three trials, each scallop was tested with (treatment) and without (control) *D. vexillum* as an epibiont. Our epibiont treatment consisted of 100% overgrowth by *D. vexillum*. To attain this level of cover, we glued, using superglue, *D. vexillum* to the top of the scallops shell, leaving enough space at the edge of the shell that cover by *D. vexillum* would not physically inhibit swimming. Shell valves were not covered. To ensure our treatment only reflected movement limited by *D. vexillum*, a similar amount of superglue was also applied to the top shell of control scallops.

To elicit an escape response, scallops were induced to swim using a seastar (*Asterias rubens*) attached, using nylon filament, to the end of a stick. At the beginning of each trial, the tubefeet of the seastar touched the scallop, thereby inducing an escape response (swimming). To examine time to initial exhaustion, the seastar continued to touch the scallop (and hence the scallop continued to swim) until it rested and further clapping was not observed, at which time the trial came to an end.

Time to initial exhaustion, swimming speed, horizontal and vertical distances swum were collected after each trial from the video footage. Initial time to exhaustion was measured as the total amount of time the scallop spent swimming. Horizontal distance was the total distance traveled while swimming. Vertical distance was measured as the greatest height in the water column reached by each scallop. Speed was

Figure 1. Individual mean measurements of total time swimming (seconds) until exhaustion as a function of shell height of sea scallop, *Placopecten magellanicus*. Gray (control) and black (treatment) dots indicate individual mean values for control (gray; no epibiosis) and treatment (black; *Didemnum vexillum* epibiosis) trials, respectively. Gray and black lines indicate regressions fitted for each of the control and treatment data sets.





Figure 2. Horizontal distance traveled by sea scallops (*Placopecten magellanicus*) as a function of shell height. Gray (control) and black (treatment) dots indicate individual mean values for control (no epibiosis) and treatment (*Didemnum vexillum* epibiosis) trials, respectively. Gray and black lines indicate regressions fitted for each of the control and treatment data sets.



calculated as the sum of the horizontal and vertical distance divided by total time swimming. This gave an average speed for that trial. Scallops (control and treatment) were made to swim to exhaustion once a day and then allowed to rest 11–14 hours, a suitable period of rest for scallops (Donovan et al. 2002). A paired-t test was used to test for differences between treatments for mean response variables.

Results

Overall there was a significant negative effect of epibiosis on the escape response of *Placopecten magellanicus* (Table 1). Sea scallops encrusted by *Didemnum vexillum* reached exhaustion more rapidly than unencrusted sea scallops (Figure 1).

They did not swim as far or as high as unencrusted *P. magellanicus* (Figures 2 and 3). Finally, encrusted sea scallops did not swim as fast as individuals free of encrustation (Figure 4).

A difference in swimming behavior was observed between encrusted and unencrusted sea scallops, however the magnitude of the difference for time to initial exhaustion and horizontal distance traveled appeared to be dependent on size (Figures 1 and 2). For time to initial exhaustion, difference between control and treatments decreased with increasing shell height (Figure 1). Smaller scallops were quicker to fatigue than larger scallops. In contrast, the divergence in distance traveled between encrusted and unencrusted sea scallops was greater for larger scallops (Figure 2).

Figure 3. Vertical distance traveled by scallops as a function of shell height. Gray (control) and black (treatment) dots indicate individual mean values for control (no epibiosis) and treatment (*Didemnum vexillum* epibiosis) trials, respectively. Gray and black lines indicate regressions fitted for each of the control and treatment data sets.




Figure 4. Swimming speed of sea scallops as a function of shell height. Gray (control) and black (treatment) dots indicate individual mean values for control (no epibiosis) and treatment (*Didemnum vexillum* epibiosis) trials, respectively. Gray and black lines indicate regressions fitted for each of the control and treatment data sets.




Table 1. Paired t-test on mean response variables (time swimming, horizontal and vertical distance) before and after treatment.

	T-statistic	Degrees of Freedom	P-value
Time to exhaustion	3.13	38	<0.003
Horizontal distance	4.51	38	<0.001
Vertical distance	2.40	38	<0.022
Swimming speed	2.72	38	<0.001

Discussion

Didemnum vexillum was first noted in Damariscotta, ME, before 1982 by oyster farmers (R. Clime, pers. comm.) and has since been observed in sea scallop habitat along the coast of Maine, NH and the Isles of Shoals (Dijkstra and L.G. Harris, pers. obs.). The sea scallop uses swimming to access a habitat with more resources (e.g., food) and to avoid predators. Our study demonstrates that encrustation by the invasive colonial ascidian *D. vexillum* negatively affects the escape response of sea scallops. Encrusted sea scallops swam less and reached exhaustion sooner than unencrusted sea scallops, specifically for scallops with shell heights of less than 40 mm.

The metabolism of smaller individuals is faster than larger individuals and likely exacerbated the time to initial exhaustion observed between control and treatment scallops.

Distance and height traveled was substantially different between encrusted and unencrusted sea scallops. Unencrusted scallops traveled farther (horizontally and vertically) than encrusted scallops. Controls swam on average 20 cm farther during each trial than treatment scallops, and the difference was greater for scallops > 40 mm. Weight of the ascidian may be proportionately greater for larger scallops than for smaller scallops. Therefore, larger scallops with heavier fouling loads may suffer greater reductions in swimming ability than smaller scallops. Reduction in the ability of encrusted *P. magellanicus* to swim as fast, as high or as far as unencrusted individuals likely increases their vulnerability to predation and reduces their ability to disperse.

It remains unclear whether scallops gain benefits from *D. vexillum* encrustation. In similar studies examining epibiosis by sponges on scallops, it was found that scallops benefited from the presence of the sponge because it was harder for seastar tube feet to grip overgrown scallop shells (Burns and Bingham 2002). *D. vexillum* may provide a similar defense. In addition, the tunic of *D. vexillum* contains acids (Bullard et al. 2007), which could further deter predation by seastars and fishes (Pisut and Pawlik 2002).

In New England, the sea scallop fishery is worth about \$431 million annually (Athearn 2005). This fishery, however, may be threatened by the continuing spread of *Didemnum vexillum* into sea scallop habitat. Unlike most invasive colonial ascidians, *D. vexillum* grows well in deep water (~49 m) and cooler habitats (Valentine et al. 2007; Valentine et al. 2009). In the Gulf of Maine, the species was found associated with scallop beds around the Isles of Shoals, ME and NH, Portsmouth Harbor, NH, Damariscotta River, ME, Georges Bank and Eastport, ME (Valentine et al. 2007; Dijkstra et al. 2007; Dijkstra and Harris, unpub. data). While the presence of *D. vexillum* is likely to inhibit recruitment of scallops in these habitats (Morris et al. 2009), growth or reproduction may also be affected by the overgrowth of *D. vexillum*, by preventing dispersal into food rich habitats. Further research on the effects of overgrowth by *D. vexillum* is needed to assess the full ecological impact of this species on sea scallops.

Acknowledgements

This study was partially funded by the Bruce McGinnis Foundation and the Shoals Marine Laboratory Research Internships in Field Science program. We would like to thank Linda Auker and the staff at the Shoals Marine Laboratory, particularly Robin Hadlock Seeley and the Research Interns in Field Science for their encouragement and assistance in field work. We also wish to thank Dana Morse, Will Hopkins and Chris Bartlett for collecting scallops.

References

- Athearn K (2005) Cobscook Bay sea scallops: the fishery and markets. Cobscook Bay Resource Center, Eastport, ME, USA, 66 pp
- Baird FT (1954) Migration of the deep sea scallop (*Pecten magellanicus*). Maine Department of Sea Shore Fisheries Circ. 14, 8 pp
- Bullard SG, Lambert G, Carman MR, Byrnes J, Whitlatch RB, Ruiz GM, Miller RJ, Harris LG, Valentine PC, Collie JS, Pederson J, McNaught DC, Cohen AN, Asch RG, Dijkstra J, Heinonen K (2007) The invasive colonial ascidian *Didemnum* sp.: current distribution, basic biology, and potential threat to marine communities of the northeast and west coasts of the United States. *Journal of Experimental Marine Biology and Ecology* 342: 99–108, <http://dx.doi.org/10.1016/j.jembe.2006.10.020>
- Burns DO, Bingham BL (2002) Epibiotic sponges on the scallops *Chlamys hastata* and *Chlamys rubida*: increased survival in a high-sediment environment. *Journal of the Marine Biological Association of the United Kingdom* 82: 961–966, <http://dx.doi.org/10.1017/S0025315402006458>
- Dijkstra J, Harris LG, Westerman E (2007) The distribution and long-term temporal patterns of four invasive colonial ascidians in the Gulf of Maine. *Journal of Experimental Marine Biology and Ecology* 342: 61–68, <http://dx.doi.org/10.1016/j.jembe.2006.10.015>
- Donovan DA, Bingham BL, Farren HM, Gallardo R, Vigilante VL (2002) Effects of sponge encrustation on the swimming behaviour energetics and morphometry of the scallop *Chlamys hastata*. *Journal of Marine Biology Association of the UK* 82: 469–476, <http://dx.doi.org/10.1017/S0025315402005738>
- Donovan DA, Bingham BL, From M, Fleisch AF, Loomis ES (2003) Effects of barnacle encrustation on the swimming behaviour, energetics, morphometry, and drag coefficient of the scallop *Chlamys hastata*. *Journal of Marine Biology Association of the UK* 89: 813–819, <http://dx.doi.org/10.1017/S0025315403007847h>
- Forester AJ (1979) The association between the sponge *Halichondria panicea* (Pallas) and scallop *Chlamys varia* (L.): A commensal protective mutualism. *Journal of Experimental Marine Biology and Ecology* 36: 1–10, [http://dx.doi.org/10.1016/0022-0981\(79\)90096-0](http://dx.doi.org/10.1016/0022-0981(79)90096-0)
- Hamilton PV, Koch KM (1996) Orientation toward natural and artificial grassbeds by swimming bay scallops, *Argopecten irradians* (Lamarck, 1819). *Journal of Experimental and Marine Biology* 199: 79–88, [http://dx.doi.org/10.1016/0022-0981\(95\)00191-3](http://dx.doi.org/10.1016/0022-0981(95)00191-3)
- Hart DR, Chute AS (2004) Essential fish habitat source document: Sea scallop, *Placopecten magellanicus*, life-history and habitat characteristics. 2nd edition NOAA Technical Memo NMFS NE–189, 21 pp

- Hart DR, Rago PJ (2006) Long-term dynamics of U.S. sea scallop (*Placopecten magellanicus*) populations. *North American Journal of Fisheries Management* 26: 490–501, <http://dx.doi.org/10.1577/M04-116.1>
- Mercer JM, Whitlatch RB, Osman RW (2009) Potential effects of the invasive colonial ascidian (*Didemnum vexillum* Kott, 2002) on pebble-cobble bottom habitats in Long Island Sound, USA. *Aquatic Invasions* 4: 133–142, <http://dx.doi.org/10.3391/ai.2009.4.1.14>
- Moore JK, Marshall N (1967) An analysis of the movements of the bay scallop, *Aequipecten irradians*, in a shallow estuary. *Proceedings of the National Shellfisheries Association* 57: 77–82
- Morris JAJ, Carman MR, Hoagland KE, Green-Beach ERM, Karney RC (2009) Impact of the invasive colonial tunicate *Didemnum vexillum* on the recruitment of the bay scallop (*Argopecten irradians irradians*) and implications for recruitment of the sea scallop (*Placopecten magellanicus*) on Georges Bank. *Aquatic Invasions* 4: 207–211, <http://dx.doi.org/10.3391/ai.2009.4.1.21>
- Morton B (1980) Swimming in *Amusium pleuronectes* (Bivalvia: Pectinidae). *Journal of Zoology* 190: 375–404, <http://dx.doi.org/10.1111/j.1469-7998.1980.tb01434.x>
- NMFS (National Marine Fisheries Service) (2007) Annual landings query: Annual commercial landings statistics. <https://www.st.nmfs.noaa.gov/st1/commercial/landings/annualandings.html> (Accessed on March 20, 2010)
- Neves CS, Rocha RM, Pitombo FB, Roper JJ (2007) Use of artificial substrata by introduced and cryptogenic marine species in Paranaguá Bay, southern Brazil. *Biofouling* 23: 319–330, <http://dx.doi.org/10.1080/08927010701399174>
- Peterson CH, Ambrose WJ, Hunt JH (1982) A field test of the swimming response of the bay scallop (*Argopecten irradians*) to changing biological factors. *Bulletin of Marine Science* 32: 939–944
- Pisut DP, Pawlik JR (2002) Anti-predatory chemical defenses of ascidians: secondary metabolites and inorganic acids? *Journal of Experimental Marine Biology and Ecology* 270: 203–214, [http://dx.doi.org/10.1016/S0022-0981\(02\)00023-0](http://dx.doi.org/10.1016/S0022-0981(02)00023-0)
- Pogsay JA (1957) The range of the sea scallop. *Nautilus* 71: 55–57
- Ruiz GM, Carlton JT, Grosholz ED, Hines AH (1997) Global invasions of marine and estuarine habitats by non-indigenous species: Mechanisms, extent, and consequences. *American Zoologist* 37: 621–632
- Stephens PJ, Boyle PR (1978) Escape responses of the queen conch *Chlamys opercularis* (L.) (Mollusca: Bivalvia). *Marine Behaviour and Physiology* 5: 103–113, <http://dx.doi.org/10.1080/10236247809378527>
- Valentine PC, Carman MR, Dijkstra J, Blackwood DS (2009) Larval recruitment of the invasive colonial ascidian *Didemnum vexillum*, seasonal water temperatures in New England coastal and offshore waters, and implications for spread of the species. *Aquatic Invasions* 4: 153–168, <http://dx.doi.org/10.3391/ai.2009.4.1.16>
- Valentine PC, Collie JS, Reid RN, Asch RG, Guida VG, Blackwood DS (2007) The occurrence of the colonial ascidian *Didemnum* sp. on Georges Bank gravel habitat: Ecological observations and potential effects on groundfish and scallop fisheries. *Journal of Experimental Marine Biology and Ecology* 342: 179–181, <http://dx.doi.org/10.1016/j.jembe.2006.10.038>