

Short Communication

Managing invasive *Styela clava* populations: Inhibiting larval recruitment with medetomidine

Kate J. Willis* and Chris M.C. Woods

National Institute of Water and Atmospheric Research (NIWA), P.O. Box 8602, Riccarton, Christchurch, New Zealand

E-mail: kjwillis68@gmail.com (KJW), c.woods@niwa.co.nz (CMCW)

*Corresponding author

Received: 3 September 2010 / Accepted 4 January 2011 / Published online: 16 June 2011

Editor's note:

This paper is a contribution to the proceedings of the 3rd International Invasive Sea Squirt Conference held in Woods Hole, Massachusetts, USA, on 26–28 April 2010. The conference provided a venue for the exchange of information on the biogeography, ecology, genetics, impacts, risk assessment and management of invasive tunicates worldwide.

Abstract

The toxicity of the synthetic catemine medetomidine to <12 h-old larval clubbed tunicate, *Styela clava*, was assessed in 2-h laboratory bioassays. Exposure to medetomidine resulted in increasing rates of larval immobility with increasing concentration. The 2-h EC50 was 3.8 mg/L medetomidine. Larval settlement was highest at a concentration of 0.1 mg/L medetomidine. At higher concentrations of medetomidine, metamorphosis was initiated but not all larvae settled. The ability of medetomidine to reduce *S. clava* larval mobility and interfere with settlement suggests that it has potential as a management tool for controlling subtidal invasive *S. clava* populations.

Key words: tunicate, population control, Selektepe™, toxicity bioassays, settlement inhibition, metamorphosis, biofouling

Globally, invasive ascidians are a growing problem, with continuing range expansion of numerous species of solitary and colonial ascidians (Lambert 2007). The Asian clubbed tunicate, *Styela clava* (Herdman, 1881), is a large (up to 20 cm total length, Minchin et al. 2006), stalked, solitary ascidian that is native to the coastal waters of the northwest Pacific (Millar 1960). This ascidian is widespread on artificial substrata in harbours, marinas, and on aquaculture infrastructure, in both hemispheres (Clarke and Therriault 2007; Nunn and Minchin 2009). In some areas, *S. clava* is also colonizing natural hard surfaces, such as crevices on rocky intertidal platforms, and shell fragments on subtidal mud flats (Goldstien et al. 2010), posing a potential threat to valued marine environments and resources, as well as aquaculture industries.

Control of solitary ascidians has focused on removal of adults from artificial structures and mussel-farming seed stock using methods such

as plucking by divers, high pressure water blasting of removable fouled structures on land, air drying, immersion in freshwater, acetic acid and chlorine exposure, encapsulation, and perforation (Carver et al. 2003; Coutts and Forrest 2005; Morrisey et al. 2008; C. Paetzold and J. Davidson pers. comm.). The development of control methods that focus on key stages in the ascidian life cycle, such as larval settlement and metamorphosis, offers alternative methods for management of invasive populations. Many natural and synthetic compounds inhibit settlement of larval ascidians (e.g. Pawlik and Hadfield 1990; Davis et al. 1991; Bellas 2006), and may provide tools that impede population growth. Medetomidine, which acts as a α_2 -adrenoceptor agonist in vertebrates (Virtanen et al. 1988), inhibits barnacle cyprid settlement at non-lethal nanomolar concentrations (Dahlström et al. 2000). This compound is being developed as a novel efficacious and environmentally




Figure 1. Mobility of *Styela clava* larvae following 2-h exposure to medetomidine in January.

sustainable antifouling agent with the brand name Selektope™, by I-Tech in Sweden (<http://www.i-tech.se>). We present the results of toxicity bioassays to determine whether medetomidine inhibits *S. clava* larval settlement, and assess the potential of this compound for management of invasive *S. clava* populations.

Adult *S. clava* were collected from a non-native population in the Port of Lyttelton (43° 36'S, 172°43'E), New Zealand, in January and March 2010 when seawater temperatures were ca. 18°C. In the laboratory, adults were maintained in large aquaria (80 L) with aerated seawater (salinity 32–34), at 18°C ($\pm 1^\circ\text{C}$), under dim fluorescent light (photoperiod 14 L: 12 D, $9.99 \mu\text{mol m}^{-2} \text{s}^{-1}$). Within 48 h of collection, up to ten adults were dissected, and gametes obtained directly from the gonoducts with a Pasteur pipette. Oocyte and sperm suspensions were pooled, mixed, and diluted with filtered (38 μm) seawater, and incubated at 18°C ($\pm 1^\circ\text{C}$) overnight with constant aeration. The following morning, only larvae actively swimming close to the water surface were collected for use in toxicity bioassays.


The acute toxicity of medetomidine to larval *S. clava* was investigated in 2-h static toxicity bioassays using <12 h-old larvae. A 99% pure formulation of medetomidine hydrochloride (Tocris Bioscience, Bristol, UK) was used to prepare a stock solution (40 mg/L) in filtered (38 μm) seawater. Larvae produced by adults collected in January were exposed to four nominal toxicant concentrations (1, 10, 20 and

40 mg/L) and a seawater control, in 20 mL glass vials containing 15 mL of test solution. For each treatment there were three replicates of ca. 20 larvae. Bioassays were undertaken in a temperature-controlled room (18°C $\pm 1^\circ\text{C}$) under dim fluorescent light. The test endpoint was larval immobility after 2 h, identified by a lack of movement when gently prodded. The median effective concentration (EC50) resulting in immobility of 50% of larvae was calculated according to the probit method (Finney 1971). Bioassays were considered successful if control survival was $\geq 90\%$ (USEPA 1991). In the bioassay with larvae from adults collected in March, the number of immobile, metamorphosed (larvae without a tail), and settled larvae were recorded following 2-h exposure to five nominal toxicant concentrations (0.1, 1, 10, 20 and 40 mg/L) and a seawater control. Due to limited numbers of larvae there were only two replicates per treatment, giving totals of 44 to 85 larvae per concentration. The results of the March bioassay are presented as means of the two replicates and EC50 values could not be calculated.

In January, exposure of <12 h-old *S. clava* larvae to medetomidine for 2 h resulted in increasing rates of larval immobility with increasing concentration compared to the control, with 20% (S.E.: 2.9%) of larvae immobile at 1 mg/L and 96% (S.E.: 4.2%) immobile at 20 mg/L medetomidine (Figure 1). The 2-h EC50 was 3.8 (95% C.I.: 1.3, 10.6) mg/L medetomidine. In the March bioassay, larvae also underwent metamorphosis and settlement, consequently numbers of mobile and immobile larvae were lower than in January. The number of settled larvae was variable, with no apparent dose-dependent response and maximum settlement (80%) occurring at 0.1 mg/L medetomidine (Figure 2). At 20 mg/L, 30% of larvae settled, while metamorphosis was initiated in 65% of larvae, but they had not settled.

Short term exposure to medetomidine reduced mobility of *S. clava* larvae and interfered with settlement at concentrations similar to those causing mortality in the barnacle *Balanus improvisus* (Dahlström et al. 2000). Larval responses to medetomidine varied between bioassays, with higher rates of metamorphosis and settlement in bioassays undertaken in late summer (March). This may be due to seasonal variations in larval competency and their propensity to settle, possibly due to differences in adult fitness at the time of spawning. In March, settlement of *S. clava* larvae during the

Figure 2. Mobility, metamorphosis, and settlement of *Styela clava* larvae following 2-h exposure to medetomidine in March.



2-h exposure was highest (80%) at 0.1 mg/L medetomidine. While metamorphosis was initiated at higher concentrations, not all larvae completed the settlement process. To determine the long-term viability of post-settlement stages, future bioassays should also monitor development through to complete metamorphosis. Medetomidine inhibits barnacle cyprid settlement by selectively binding to octopamine receptors, causing the legs to become hyperactive and unable to attach to surfaces (Lind et al. 2010). The mode of action of medetomidine is unknown in ascidians, but its ability to reduce *S. clava* larval mobility and interfere with settlement, suggests that it has potential as a management tool for controlling subtidal invasive populations by reducing recruitment rates, and ultimately population growth. Given a planktonic larval phase of <24 h (Davis 1997), and the difficulty of ensuring extended exposure in the marine environment, our 2 h exposures attempted to reproduce possible treatment scenarios. The application of mitigation compounds, such as medetomidine, that target ascidian early developmental stages necessitates knowledge of local reproductive and spawning behaviour, and development of underwater application techniques, such as antifouling coatings, slow-release pellets, gel adjuvants, and microencapsulated beads, to enable topical application and extended exposure during peak spawning activity.

Acknowledgements

Thanks to Lisa Peacock, Kate Schimanski, and Caroline Williams for collection of adult *Styela clava*, and two anonymous reviewers for their comments on the manuscript. Access to structures within the Port of Lyttelton was granted by the Lyttelton Port Company Ltd. This work was funded by NIWA Capability Fund CF105044.

References

- Bellas J (2006) Comparative toxicity of alternative antifouling biocides on embryos and larvae of marine invertebrates. *Science of the Total Environment* 367: 573–585, <http://dx.doi.org/10.1016/j.scitotenv.2006.01.028>
- Carver CE, Chisholm A, Mallet AL (2003) Strategies to mitigate the impact of *Ciona intestinalis* (L.) biofouling on shellfish production. *Journal of Shellfish Research* 22: 621–631
- Clarke CL, Theriault TW (2007) Biological synopsis of the invasive tunicate *Styela clava* (Herdman 1881). Canadian Manuscript Report of Fisheries and Aquatic Sciences 2807, vi+23 pp
- Coutts ADM, Forrest BM (2005) Evaluation of eradication tools for the clubbed tunicate *Styela clava*. Cawthron Report, vol. 1110. Cawthron Institute, Nelson, New Zealand. December 2005, 48 pp
- Dahlström M, Mårtensson LGE, Jonsson PR, Arnebrant T, Elwing H (2000) Surface active adrenoceptor compounds prevent the settlement of cyprid larvae of *Balanus improvisus*. *Biofouling* 16: 191–203, <http://dx.doi.org/10.1080/08927010009378444>
- Davis AR, Butler AJ, van Altena I (1991) Settlement behaviour of ascidian larvae: preliminary evidence for inhibition by sponge allelochemicals. *Marine Ecology Progress Series* 72: 117–123, <http://dx.doi.org/10.3354/meps072117>

- Davis MH (1997) Physical factors influencing larval behaviour in three species of solitary ascidian. PhD thesis, Open University. 360 pp
- Finney DJ (1971) Probit analysis. Cambridge University Press, 333 pp
- Goldstien SJ, Schiel DR, Gemmell NJ (2010) Regional connectivity and coastal expansion: differentiating pre-border and post-border vectors for the invasive tunicate *Styela clava*. *Molecular Ecology* 19: 874–885, <http://dx.doi.org/10.1111/j.1365-294X.2010.04527.x>
- Lambert G (2007) Invasive sea squirts: A growing global problem. *Journal of Experimental Marine Biology and Ecology* 342: 3–4, <http://dx.doi.org/10.1016/j.jembe.2006.10.009>
- Lind U, Rosenblad MA, Frank LH, Falkbring S, Brive L, Jonne M, Laurila JM, Pohjanoksa K, Vuorenpää A, Kukkonen JP, Gunnarsson L, Scheinin M, Mårtensson Lindblad LGE, Blomberg A (2010) Octopamine receptors from the barnacle *Balanus improvisus* are activated by the α_2 -adrenoceptor agonist medetomidine. *Molecular Pharmacology* 78: 237–248, <http://dx.doi.org/10.1124/mol.110.063594>
- Millar H (1960) The identity of the ascidians *Styela mammiculata* Carlisle and *S. clava* Herdman. *Journal of the Marine Biological Association of the United Kingdom* 39: 509–511, <http://dx.doi.org/10.1017/S0025315400013503>
- Minchin D, Davis MH, Davis ME (2006) Spread of the Asian tunicate *Styela clava* Herdman, 1882 to the east and south-west coasts of Ireland. *Aquatic Invasions* 1: 91–96, <http://dx.doi.org/10.3391/ai.2006.1.2.7>
- Morrisey D, Page M, Handley S, Middleton C, Schick R (2008) Biology and ecology of the introduced ascidian *Eudistoma elongatum*, and trials of potential methods for its control. MAF Biosecurity New Zealand Technical Paper. May 2008
- Nunn JD, Minchin D (2009) Further expansions of the Asian tunicate *Styela clava* Herdman, 1882 in Ireland. *Aquatic Invasions* 4: 591–596, <http://dx.doi.org/10.3391/ai.2009.4.4.4>
- Pawlik JR, Hadfield MG (1990) A symposium on chemical factors that influence this settlement and metamorphosis of marine invertebrate larvae: introduction and perspective. *Bulletin of Marine Science* 46: 450–454
- USEPA (United States Environmental Protection Agency) (1991) Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms. Cincinnati, Ohio
- Virtanen R, Savola JM, Saano V, Nyman L (1988) Characterisation of the selectivity, specificity and potency of medetomidine as an α_2 -adrenoceptor agonist. *European Journal of Pharmacology* 150: 9–14, [http://dx.doi.org/10.1016/0014-2999\(88\)90744-3](http://dx.doi.org/10.1016/0014-2999(88)90744-3)